

Active Chitinase-3-like Protein 1 (CHI3L1) Instruction Manual

SBPB230Hu01

Homo sapiens (Human)

Buffer Formulation

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits

Freeze-dried powder

Purity

> 97%

Isoelectric Point

9.4

Applications

Cell culture; Activity Assays.

ACTIVITY TEST




Figure. Cell proliferation of Raji cells after stimulated with GP39.




Figure. Cell proliferation of Raji cells after stimulated with GP39. Glycoprotein 39, Cartilage (GP39) also known as Chitinase-3-like protein 1 (CHI3L1) is a secreted glycoprotein that is approximately 40kDa in size that in humans is encoded by the CHI3L1 gene. GP39 plays a role in cancer cell proliferation, survival, invasiveness and in the regulation of cell-matrix interactions. To test the effect of GP39 on cell proliferation, Raji cells were seeded into triplicate wells of 96-well plates at a density of 5,000 cells/well with 1% serum standard 1640 which contains various concentrations of recombinant human GP39. After incubated for 5 days, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 μ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of Raji cells after incubation with GP39 for 5 days observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant GP39 for 5 days. The result was shown in Figure 2. It was obvious that GP39 significantly increased cell viability of Raji cells.

(A) Raji cells cultured in 1640, stimulated with 10ng/mL GP39 for 5 days;

(B) Unstimulated Raji cells cultured in 1640 for 5 days.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE




Figure. Western Blot

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.