

Active Endothelial Cell Specific Molecule 1 (ESM1) Instruction Manual

SBPC230Hu01

Homo sapiens (Human)

Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.7
Applications	Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of ESM1 with LFA1a.

Endothelial cell-specific molecule 1 (ESM1) is a proteoglycan secreted by endothelial cells (primarily in the human lung and kidney tissues) and its mRNA expression is regulated by inflammatory cytokines. Endocan expression, which detected in various epithelia and adipocytes has been shown to be upregulated by vascular endothelial growth factor (VEGF), fibroblast growth factor 2 (FGF-2), TNF alpha, IL1 beta, or lipopolysaccharide and downregulated by IFN gamma. Genetically engineered cells overexpressing ESM1 induce tumor formation, implying that ESM1 might be involved in the pathophysiology of tumor growth in vivo. Besides, Lymphocyte Function Associated

Antigen 1 Alpha (LFA1a) has been identified as an interactor of ESM1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ESM1 and recombinant human LFA1a. Briefly, ESM1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to LFA1a-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti- ESM1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of of ESM1 and LFA1a was shown in Figure 1, and this effect was in a dose dependent manner.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image




Figure. Western Blot; Sample: Recombinant ESM1, Human.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.