

Eukaryotic Cluster Of Differentiation 147 (CD147) Instruction Manual

SFPB438Ra61

Rattus norvegicus (Rat)

Source	Eukaryotic expression
Host	293F cell
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	34.6kDa
Accurate Molecular Mass	44-55kDa(Analysis of differences refer to the manual)
Residues & Tags	Ala22~Arg324 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	4.9
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
                AAGFLKAPM  SQEQWAGGSV  VLHCEAVGSP  
MPEIQWVFEG  NEPNDSCSQL  WDGARLDRVH  IHATYRQHAA  STLSVDGLAA  
EDTGTYECRA  SSDPDRNHLT  RPPRVKQVRA  QASVVVLEPG  TIVTSVQEVD  
SKTQLTCFLN  SSGIDIVGHR  WMRGGKVLQE  DTLPLDQMKY  TVDADDRSGE  
YSCIFLPEPV  GRGNINVEGP  PRIKVGKKSE  HASEGEFVKL  ICKSEASHPP  
VDEWVFKTS  DTGDQTISNG  TEANSKYVII  STPELSELII  SLDLMMNVDPG  
TYVCNATNSQ  GSARETISLR  VRSR
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.