



Rabbit Anti-TGF Beta 1 antibody

SL0103R

Product Name:	TGF Beta 1
Chinese Name:	转化生长因子β1抗体
Alias:	CED; DPD1; LAP; Latency-associated peptide; TGF beta 1; TGF beta; TGF-beta 1; TGF beta 1 protein; TGF-beta 1 protein; TGF-beta-1; TGF-beta-5; TGF-beta1; TGFB; Tgfb-1; tgb1; TGFB1_HUMAN; TGFbeta; TGFbeta1; Transforming Growth Factor b1; Transforming Growth Factor beta 1; Transforming growth factor beta 1a; transforming growth factor beta-1; transforming growth factor, beta 1.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	44kDa
Cellular localization:	Extracellular matrixSecretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human TGF beta 1:101-150/390
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types. Many cells have TGFB

receptors, and the protein positively and negatively regulates many other growth factors. The secreted protein is cleaved into a latency-associated peptide (LAP) and a mature TGFB1 peptide, and is found in either a latent form composed of a TGFB1 homodimer, a LAP homodimer, and a latent TGFB1-binding protein, or in an active form composed of a TGFB1 homodimer. The mature peptide may also form heterodimers with other TGFB family members. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.

Function:

Multifunctional protein that controls proliferation, differentiation and other functions in many cell types. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other growth factors. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts.

Subunit:

Homodimer; disulfide-linked, or heterodimer with TGFB2. Secreted and stored as a biologically inactive form in the extracellular matrix in a 290 kDa complex (large latent TGF-beta1 complex) containing the TGFB1 homodimer, the latency-associated peptide (LAP), and the latent TGFB1 binding protein-1 (LTBP1). The complex without LTBP1 is known as the 'small latent TGF-beta1 complex'. Dissociation of the TGFB1 from LAP is required for growth factor activation and biological activity. Release of the large latent TGF-beta1 complex from the extracellular matrix is carried out by the matrix metalloproteinase MMP3 (By similarity). May interact with THSD4; this interaction may lead to sequestration by FBN1 microfibril assembly and attenuation of TGFB signaling. Interacts with the serine proteases, HTRA1 and HTRA3: the interaction with either inhibits TGFB1-mediated signaling. The HTRA protease activity is required for this inhibition (By similarity). Interacts with CD109, DPT and ASPN.

Subcellular Location:

Secreted, extracellular space, extracellular matrix.

Tissue Specificity:

Highly expressed in bone. Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA). Co-localizes with ASPN in chondrocytes within OA lesions of articular cartilage.

Post-translational modifications:

Glycosylated.

The precursor is cleaved into mature TGF-beta-1 and LAP, which remains non-covalently linked to mature TGF-beta-1 rendering it inactive.

DISEASE:

Defects in TGFB1 are the cause of Camurati-Engelmann disease (CE) [MIM:131300]; also known as progressive diaphyseal dysplasia 1 (DPD1). CE is an autosomal dominant disorder characterized by hyperostosis and sclerosis of the diaphyses of long bones. The

disease typically presents in early childhood with pain, muscular weakness and waddling gait, and in some cases other features such as exophthalmos, facial paralysis, hearing difficulties and loss of vision.

Similarity:

Belongs to the TGF-beta family.

SWISS:

P01137

Gene ID:

7040

Database links:

[Entrez Gene: 7040](#)Human

[Entrez Gene: 21803](#)Mouse

[Entrez Gene: 59086](#)Rat

[Omim: 190180](#)Human

[SwissProt: P01137](#)Human

[SwissProt: P04202](#)Mouse

[SwissProt: P17246](#)Rat

[Unigene: 645227](#)Human

[Unigene: 248380](#)Mouse

[Unigene: 40136](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Growth factors and hormones (Growth Factor and Hormones)


TGF是一种多效生长因子, 对epithelial cells增值有潜在抑制作用, 可抑制Tumour生长用于许多恶性Tumour如:胃癌、肺癌、膀胱癌、肾癌、前列腺癌、结肠癌等多种恶性Tumour的研究。

TGF β 超级家族由为数众多的生长的分化因子组成,包括转移生长因子 β 1, 2和3(TGF β 1, TGF β 2, TGF β 3);胎盘生长因子(PTGF- β);生长/分化因子(GDFs);缪氏抑制物(MIS);骨形态形成蛋白(BMPs);交织细胞元神经生长因子(GDNF);抑制素和活化素(α , β -A,和 β -C), Lefty和Nodal。

TGF超级家族成员参与胚胎发育和成体组织体内平衡。TGF- β 1抗体与TGF-

β2和TGF-β3没有React Species。与豚鼠有大部分交叉。

Picture:




Sample: Lymph node(Mouse) Lysate at 30 ug

Primary: Anti- TGF Beta 1(SL0103R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution

Predicted band size: 44 kD

Observed band size: 54 kD




Sample: Bones(Mouse) Lysate at 30 ug

Primary: Anti- TGF Beta 1(SL0103R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution

Predicted band size: 44 kD

Observed band size: 55 kD




Sample: K562 Cell Lysate at 40 ug

Primary: Anti- TGF Beta 1(SL0103R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 44 kD

Observed band size: 60 kD




bs-0103R Anti-TGF-beta1

Formalin-fixed and paraffin-embedded human Pituitary tumor labeled with Rabbit Anti- TGF-beta1 Polyclonal Antibody, Unconjugated(bs-0103R) at 1:300 followed by conjugation to the secondary antibody and DAB staining

Tissue/cell: human Pituitary tumor; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-TGF-beta1 Polyclonal Antibody, Unconjugated(SL0103R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Oral squamous cell carcinoma;4% Paraformaldehyde-fixed;

Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-TGF-beta1 Polyclonal Antibody, Unconjugated(SL0103R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, PE-Cy3 conjugated(SL0103R)used at 1:200 dilution for 40 minutes at 37°C.

Excitation wavelength: 530nm; Emission wavelength:578nm