



## Rabbit Anti-p75 NGF Receptor antibody

SL0161R

<b>Product Name:</b>	p75 NGF Receptor
<b>Chinese Name:</b>	神经生长因子受体抗体
<b>Alias:</b>	NGFR; Gp80 LNGFR; ngfrp75; Low affinity nerve growth factor receptor; Low affinity neurotrophin receptor p75NTR; NGF receptor; p75 ICD; CD271; CD271 antigen; Gp80 LNGFR; Low affinity nerve growth factor receptor; Low affinity neurotrophin receptor p75NTR; Nerve growth factor receptor; Nerve growth factor receptor TNFR superfamily member 16; NGF receptor; NGFR; p75 ICD; p75 Neurotrophin receptor; p75 NTR; TNFR Superfamily Member 16; TNFRSF16; Tumor necrosis factor receptor superfamily member 16; TNFR16_HUMAN; NGFR p75.
<b>文献引用</b> <b>PubMed</b> :	<p><b>Specific References(3)</b> SL0161R has been referenced in 3 publications.</p> <p><b>[IF=3.31]</b>Paczkowska, Edyta, et al. "Humoral Activity of Cord Blood-Derived Stem/Progenitor Cells: Implications for Stem Cell-Based Adjuvant Therapy of Neurodegenerative Disorders." PloS one 8.12 (2013): e83833.<b>Human</b>.  <a href="#">PubMed:24391835</a></p> <p><b>[IF=1.64]</b>Shan, Tengfei, et al. "Lithium chloride promotes the odontoblast differentiation of hair follicle neural crest cells by activating Wnt/<math>\beta</math>-catenin signaling." Cell Biology International (2014).<b>Mouse</b>.  <a href="#">PubMed:25044369</a></p> <p><b>[IF=3.25]</b>?mieszek, Agnieszka, et al. "Antioxidant and Anti-Senescence Effect of Metformin on Mouse Olfactory Ensheathing Cells (mOECs) May Be Associated with Increased Brain-Derived Neurotrophic Factor Levels—An Ex Vivo Study." International Journal of Molecular Sciences 18.4 (2017): 872.<b>IF(ICC);Mouse</b>.  <a href="#">PubMed:28425952</a></p>
<b>Organism Species:</b>	Rabbit

<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human, Mouse, Rat, Dog, Pig, Cow, Horse, Guinea Pig,
<b>Applications:</b>	WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 Flow-Cyt=1µg/Test IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	44kDa
<b>Cellular localization:</b>	The nucleus cytoplasmic The cell membrane Extracellular matrix
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human NGFR:301-400/427<Cytoplasmic>
<b>Isotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	<p>The low affinity NGFR (Nerve growth factor receptor) is a 75kDa membrane-spanning glycoprotein lacking intrinsic tyrosine kinase activity. p75NGFR interacts with TrkA, the high affinity NGF receptor and potentiates TrkA signaling at low NGF concentrations. The p75 receptor binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75NGFR plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death. It is expressed on many neuronal cells types including many embryonic forms and the receptor can be used to isolate neuronal progenitor cells. NGF is important for the development, differentiation and survival of a variety of neuronal and non-neuronal cells. Its action is mediated by binding to two distinct receptors, the high affinity p140 and low affinity p75. p75NGFR binds neurotrophins including brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3), NT-4/5, and NT-6. p75NGFR belongs to the TNF-R superfamily and is reported to mediate NGF-induced apoptosis.</p> <p><b>Function:</b> Low affinity receptor which can bind to NGF, BDNF, NT-3, and NT-4. Can mediate cell survival as well as cell death of neural cells.</p> <p><b>Subunit:</b> Homodimer; disulfide-linked. Interacts with p75NTR-associated cell death executor. Interacts with NGFRAP1/BEX3. Interacts with TRAF2, TRAF4, TRAF6, PTPN13 and RANBP9. Interacts through TRAF6 with SQSTM1 which bridges NGFR to NTRK1. Interacts with BEX1. Interacts with KIDINS220 and NTRK1. Can form a ternary complex with NTRK1 and KIDINS220 and this complex is affected by the expression levels of KIDINS220. An increase in KIDINS220 expression leads to a decreased</p>

association of NGFR and NTRK1. Interacts with LINGO1 (By similarity). Interacts with NTRK2; may regulate the ligand specificity of the NTRK2 receptor. Interacts with NRADD.

**Subcellular Location:**

Membrane; Single-pass type I membrane protein.

**Post-translational modifications:**

N- and O-glycosylated.

Phosphorylated on serine residues.

**Similarity:**

Contains 1 death domain.

Contains 4 TNFR-Cys repeats.

**SWISS:**

P07174

**Gene ID:**

4804

**Database links:**

[Entrez Gene: 4804](#) Human

[Entrez Gene: 18053](#) Mouse

[Entrez Gene: 24596](#) Rat

[Omim: 162010](#) Human

[SwissProt: P08138](#) Human

[SwissProt: Q9Z0W1](#) Mouse

[SwissProt: P07174](#) Rat

[Unigene: 415768](#) Human

[Unigene: 681726](#) Human

[Unigene: 283893](#) Mouse

[Unigene: 10980](#) Rat

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


Neurobiology相关蛋白 (Neurobiology)

NGFR神经生长因子受体又称:CD271存在与细胞上的与神经生长因子相结合并相互作用的蛋白质因子, 属一型跨膜TNF受体超家族glycoprotein, 它可与NGF, BDNF, NT-3, NT-4结合, 根据与神经生长因子的结合力速度及亲和力可分为高亲和力和低亲和力两种NGFR。

此抗体识别分子量为47kDa低亲和力的NGFR.NGF-

R也称为低亲和神经生长因子受体, 属1型跨膜TNF受体超家族。

Picture:



Sample:

Brain(Rat) lysate at 30ug;


Heart (Rat)lysate at 30ug;

Primary: Anti-NGFR/p75NTR (SL0161R) at 1:200 dilution;

Secondary: HRP conjugated Goat-Anti-Rabbit IgG(bse-0295G) at 1: 3000 dilution;

Predicted band size : 44kD

Observed band size : 32kD, 44kD



Blank control (blue line): Mouse spleen (blue).

Primary Antibody (green line): Rabbit Anti- p75 NGF Receptor antibody (SL0161R)

Dilution: 1 $\mu$ g /10<sup>6</sup> cells;

Isotype Control Antibody (orange line): Rabbit IgG .


Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC

Dilution: 1 $\mu$ g /test.

Protocol

The cells were fixed with 70% ice-cold methanol overnight at 4°C and then

permeabilized with 0.1% PBS-Tween for 20 min at room temperature (The cells were fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice.). Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in 1 X PBS/2%BSA/10% goat serum to block non-specific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: (mo)nephrocyte(blue)

Isotype Control Antibody: Rabbit IgG -AF647(orange); Primary Antibody Dilution:  
5 $\mu$ l in 100  $\mu$ L 1X PBS containing 0.5% BSA(green).