

Rabbit Anti-rat IgG/AP antibody

SL0293R-AP

t Anti-rat IgG/AP
磷酸酶(AP)标记 的兔抗大鼠 IgG
noglobulin G
t
lonal
:1000-10000ELISA=1:1000-10000IHC-P=1:100-1000IHC-F=1:100-1000 t tested in other applications. al dilutions/concentrations should be determined by the end user.
Da
ilized or Liquid
ml
ength plasma protein:
N.
y purified by Protein A
TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
ge: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The ilized antibody is stable at room temperature for at least one month and for greater year when kept at -20oC. When reconstituted in sterile distilled water or diluent ed, theantibody is stable for at least two weeks at 2-4 °C.
noglobulin G (IgG), is one of the most abundant proteins in serum with normal between 8-17 mg/mL in adult blood. IgG is important for our defence against organisms and the molecules are produced by B lymphocytes as a part of our ve immune response. The IgG molecule has two separate functions; to bind to the gen that elicited the response and to recruit other cells and molecules to destroy tigen. The variability of the IgG pool is generated by somatic recombination and mber of specificities in an individual at a given time point is estimated to be 1011 ts. rtant Note:
n

Т	This product as supplied is intended for research use only, not for use in human,
tł	nerapeutic or diagnostic applications.

www.sunionobiotech.com