



## Goat Anti-rabbit IgG/Cy5 antibody

SL0295G-Cy5

<b>Product Name:</b>	Goat Anti-rabbit IgG/Cy5
<b>Chinese Name:</b>	Cy5标记的羊抗兔IgG
<b>Alias:</b>	Immunoglobulin G
<b>文献引用</b> <b>PubMed</b> :	<p><b>Specific References(3)</b> SL0295G-Cy5 has been referenced in 3 publications.</p> <p><b>[IF=5.47]</b>Pan, Bo, et al. "c-Abl Tyrosine Kinase Mediates Neurotoxic Prion Peptide-Induced Neuronal Apoptosis via Regulating Mitochondrial Homeostasis." <i>Molecular Neurobiology</i> (2014): 1-15.<b>Rat.</b>  <a href="#">PubMed:24510275</a></p> <p><b>[IF=4.46]</b>Song, Zhi-Qi, et al. "Overexpression of BAT3 Alleviates Prion Protein Fragment PrP106-126-Induced Neuronal Apoptosis." <i>CNS Neuroscience &amp; Therapeutics</i> (2014).<b>Human.</b>  <a href="#">PubMed:24629137</a></p> <p><b>[IF=4.27]</b>Dai, Wenwen, et al. "Antiviral effects of Retro-2 cycl and Retro-2.1 against Enterovirus 71 in vitro and in vivo." <i>Antiviral Research</i> (2017).<b>IF(ICC);</b>  <a href="#">PubMed:28688753</a></p>
<b>Organism Species:</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Rb
<b>Applications:</b>	Flow-Cyt=1:100-1000IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	150kDa
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	2mg/1ml

<b>immunogen:</b>	Full length plasma protein:
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent supplied, theantibody is stable for at least two weeks at 2-4 °C.
<b>Product Detail:</b>	<p>Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.</p> <p><b>Important Note:</b> This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p>

www.sunlongbiotech.com