



Goat Anti-rabbit IgG/FITC antibody

SL0295G-FITC

Product Name:	Goat Anti-rabbit IgG/FITC
Chinese Name:	FITC标记的羊抗兔IgG
Alias:	Immunoglobulin G
文献引用 :	Specific References(25) SL0295G-FITC has been referenced in 25 publications. [IF=7.60] Xu, Haixing, Yuhua Yan, and Shipu Li. "PDLLA/chondroitin sulfate/chitosan/NGF conduits for peripheral nerve regeneration." <i>Biomaterials</i> 32.20 (2011): 4506-4516. Rat. PubMed:21397324
	[IF=4.58] Lu, Cong-Xiao, et al. "Sulfated polymannuroguronate, a novel anti-AIDS drug candidate, inhibits HIV-1 Tat-induced angiogenesis in Kaposi's sarcoma cells." <i>Biochemical pharmacology</i> 74.9 (2007): 1330-1339. Human. PubMed:17868650
	[IF=3.73] Su, Bao, et al. "In Vitro and In Vivo Evaluations of Nano-Hydroxyapatite/Polyamide 66/Glass Fibre (n-HA/PA66/GF) as a Novel Bioactive Bone Screw." <i>PLOS ONE</i> 8.7 (2013): e68342. Mouse. PubMed:23861888
	[IF=3.73] Yan, Yu-xian, et al. "Mechanical strain regulates osteoblast proliferation through integrin-mediated ERK activation." <i>PLoS One</i> 7.4 (2012): e35709. Mouse. PubMed:22539993
	[IF=2.93] Liu, Yang, et al. "A Simple Method for Isolating and Culturing the Rat Brain Microvascular Endothelial Cells." <i>Microvascular Research</i> (2013). Rat. PubMed:23978334

[IF=2.38]Zhang, Feng-Lei, Hai-Qing Gao, and Lin Shen. "Inhibitory effect of GSPE on RAGE expression induced by advanced glycation end products in endothelial cells." Journal of cardiovascular pharmacology 50.4 (2007): 434-440.

[PubMed:18049312](#)

[IF=2.24]Zhang, Wan-Dang Wang, et al. "Assessing the role of IL-35 in colorectal cancer progression and prognosis." Int J Clin Exp Pathol 6.9 (2013): 1806-1816.**Human.**

[PubMed:24040445](#)

[IF=2.00]Wang, Yan, et al. "The Chronic Effects of Low Lead Level on The Expressions of Nrf2 and Mrp1 of The Testes in The Rats." Environmental Toxicology and Pharmacology (2012).**IHC-P;Rat.**

[PubMed:23274417](#)

[IF=1.73]Liu, Dong, and Zhong-Xiang Niu. "Cloning of a gene fragment encoding chicken complement component C3d with expression and immunogenicity of Newcastle disease virus F gene?CC3d fusion protein." Avian Pathology 37.5 (2008): 477-485.**iWB;Chicken.**

[PubMed:8798021](#)

[IF=1.31]Wang, Chunqiang, Wei Ma, and Yuhong Su. "NF- κ B Pathway Contributes to Cadmium-Induced Apoptosis of Porcine Granulosa Cells." Biological trace element research (2013): 1-8.**Pig.**

[PubMed:23575899](#)

[IF=0.95]Li, Fa-Rong, et al. "Reversal effect of rosmarinic acid on multidrug resistance in SGC7901/Adr cell." Journal of Asian natural products research ahead-of-print (2013): 1-10.

[PubMed:23421517](#)

[IF=5.47]Pan, Bo, et al. "c-Abl Tyrosine Kinase Mediates Neurotoxic Prion Peptide-Induced Neuronal Apoptosis via Regulating Mitochondrial Homeostasis." Molecular Neurobiology (2014): 1-15.**Rat.**

[PubMed:24510275](#)

[IF=4.46]Song, Zhi-Qi, et al. "Overexpression of BAT3 Alleviates Prion Protein Fragment PrP106-126-Induced Neuronal Apoptosis." CNS Neuroscience & Therapeutics (2014).**Human.**

[PubMed:24629137](#)

[IF=3.02]Diao, Jin-Mei, et al. "Construction of a human corneal stromal equivalent with non-transfected human corneal stromal cells and acellular porcine corneal stromata." *Experimental Eye Research* (2015).

[PubMed:25613074](#)

[IF=2.47]Chen, Han, et al. "Scorpion venom activates natural killer cells in hepatocellular carcinoma via the NKG2D-MICA pathway." *International Immunopharmacology* 35 (2016): 307-314.FCM;Chicken, Human.

[PubMed:27089390](#)

[IF=1.55]Zhang, Xumin, et al. "Phenotypic transformation of smooth muscle cells from porcine coronary arteries is associated with connexin 43." *Molecular Medicine Reports* (2016).IF(IHC-F);Pig.

[PubMed:27175888](#)

[IF=1.28]Zhang, Ying, et al. "Effects of testosterone on the expression levels of AMH, VEGF and HIF α in mouse granulosa cells." *Experimental and Therapeutic Medicine* 12.2 (2016): 883-888.IF(ICC);Mouse.

[PubMed:27446291](#)

[IF=9.11]Xia, Dan, et al. "MARCH2 regulates autophagy by promoting CFTR ubiquitination and degradation, and PIK3CA-AKT-MTOR signaling." *Autophagy* (2016).IF(ICC);

[PubMed:27308891](#)

[IF=1.26]Hu, Xiaoxiao, et al. "Ergosterol reverses multidrug resistance in SGC7901/Adr cells." *Die Pharmazie-An International Journal of Pharmaceutical Sciences*69.5 (2014): 396-400.FCM;

[PubMed:24855835](#)

[IF=5.38]Hu, Jia, et al. "TMEM166/EVA1A interacts with ATG16L1 and induces autophagosome formation and cell death." *Cell Death & Disease* 7.8 (2016): e2323.IF(ICC);Human.

[PubMed:27490928](#)

[IF=1.56]Liu, Xiao?Ya, et al. "Fibronectin expression is critical for liver fibrogenesis in vivo and in vitro." *Molecular Medicine Reports* (2016).other;All.

[PubMed:27572112](#)

	<p>[IF=2.88]Sun, Kwang-Hsiao, et al. "In vivo study of alginate hydrogel conglomerating cells to polycaprolactone vascular scaffolds fabricated by electrospinning." Journal of Biomedical Materials Research Part B: Applied Biomaterials (2016).IF(ICC); PubMed:27654960</p> <p>[IF=2.37]Zhu, Yue, et al. "Effect of TCEA3 on the differentiation of bovine skeletal muscle satellite cells." Biochemical and Biophysical Research Communications (2017).IF(ICC); PubMed:28161635</p> <p>[IF=2.57]Vuletic, Ivan, et al. "Validation of Bevacizumab Therapy Effect on Colon Cancer Subtypes by Using Whole Body Imaging in Mice." Molecular Imaging and Biology: 1-10.IF(IHC-F); PubMed:28315202</p> <p>[IF=6.64]He, Xia, et al. "Prenatal nicotine exposure induces HPA axis-hypersensitivity in offspring rats via the intrauterine programming of up-regulation of hippocampal GAD67." Archives of Toxicology (2017): 1-17.IF(IHC-F); PubMed:28555334</p>
Organism Species:	Goat
Clonality:	Polyclonal
React Species:	Rb
Applications:	Flow-Cyt=1:100-1000IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	150kDa
Form:	Lyophilized or Liquid
Concentration:	2mg/1ml
immunogen:	Full length plasma protein:
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent supplied, theantibody is stable for at least two weeks at 2-4 °C.
Product Detail:	Immunoglobulin G (IgG) is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and

the number of specificities in an individual at a given time point is estimated to be 1011 variants.

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com