



Goat Anti-human IgG/FITC antibody

SL0297G-FITC

Product Name:	Goat Anti-human IgG/FITC
Chinese Name:	FITC标记的羊抗人IgG
Alias:	Immunoglobulin G
文献引用 PubMed :	<p>Specific References(3) SL0297G-FITC has been referenced in 3 publications.</p> <p>[IF=1.79]Zhang, Ruohu, et al. "Immunoassays based on surface-enhanced fluorescence using gap-plasmon-tunable Ag bilayer nanoparticle films." Journal of fluorescence 23.1 (2013): 71-77.ELISA;Human. PubMed:22890683</p> <p>[IF=2.47]Xun, Yiping, et al. "Identification of prohibitin as an antigen in Behcet's disease." Biochemical and Biophysical Research Communications (2014).Human. PubMed:25091478</p> <p>[IF=3.53]Du H, Shi L, Chen P, Yang W, Xun Y, Yang C, et al. (2015) Prohibitin Is Involved in Patients with IgG4 Related Disease. PLoS ONE 10(5): e0125331. PubMed:25932630</p>
Organism Species:	Goat
Clonality:	Polyclonal
React Species:	hu
Applications:	Flow-Cyt=1:100-1000IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	160kDa
Form:	Lyophilized powder
Concentration:	2mg/1ml
immunogen:	Full length plasma protein:
Lsotype:	IgG

Purification:	affinity purified by Protein G
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent supplied, theantibody is stable for at least two weeks at 2-4 °C.
Product Detail:	<p>Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10¹¹ variants.</p> <p>Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p>

www.sunlongbiotech.com