

Rabbit Anti-human IgG/PE-Cy5.5 antibody

SL0297R-PE-Cy5.5

Chinese Name: PE-Cy5.5标记的兔抗人IgG Alias: Immunoglobulin G Organism Species: Rabbit Clonality: Polyclonal React Species: hu IF=1:100-1000 Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/Iml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or disapplied, theantibody is stable for at least two weeks at 2-4 °C.	
Organism Species: Rabbit Clonality: Polyclonal React Species: hu IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or	
Clonality: Polyclonal React Species: hu IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
React Species: hu IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Applications: IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
optimal dilutions/concentrations should be determined by the end user. Molecular weight: 160kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Molecular weight:160kDaForm:Lyophilized or LiquidConcentration:2mg/1mlimmunogen:Full length plasma protein:Lsotype:IgGPurification:affinity purified by Protein AStorage Buffer:0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.Storage:Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Concentration: immunogen: Full length plasma protein: Lsotype: IgG Purification: Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
immunogen: Lsotype: IgG Purification: Storage Buffer: O.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: S	
Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled water or distilled.	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled wa	
Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or	
Storage: lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled wa	
than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
than a year when kept at -200C. When reconstituted in sterile distilled water or di	
supplied, theantibody is stable for at least two weeks at 2-4 °C.	luent
Immunoglobulin G (IgG), is one of the most abundant proteins in serum with nor	
levels between 8-17 mg/mL in adult blood. IgG is important for our defence again	
microorganisms and the molecules are produced by B lymphocytes as a part of ou	
adaptive immune response. The IgG molecule has two separate functions; to bind	
Product Detail: pathogen that elicited the response and to recruit other cells and molecules to desi	
the antigen. The variability of the IgG pool is generated by somatic recombination	
the number of specificities in an individual at a given time point is estimated to be	3 1011
variants.	
Important Note:	

This product as supplied is intended for research use only, not for use in human	,
therapeutic or diagnostic applications.	

