

Rabbit Anti-human IgG/RBITC antibody

SL0297R-RBITC

| Product Name: | Rabbit Anti-human IgG/RBITC |
|-----------------------|--|
| Chinese Name: | 罗丹明标记的兔抗人IgG |
| Alias: | Immunoglobulin G |
| Organism Species: | Rabbit |
| Clonality: | Polyclonal |
| React Species: | hu |
| | IF=1:100-1000 |
| Applications: | not yet tested in other applications. |
| | optimal dilutions/concentrations should be determined by the end user. |
| Molecular weight: | 160kDa |
| Form: | Lyophilized or Liquid |
| Concentration: | 2mg/1ml |
| immunogen: | Full length plasma protein: |
| Lsotype: | IgG |
| Purification: | affinity purified by Protein A |
| Storage Buffer: | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage: | Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The |
| | lyophilized antibody is stable at room temperature for at least one month and for greater |
| | than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent |
| | supplied, theantibody is stable for at least two weeks at 2-4 °C. |
| Product Detail: | Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal |
| | levels between 8-17 mg/mL in adult blood. IgG is important for our defence against |
| | microorganisms and the molecules are produced by B lymphocytes as a part of our |
| | adaptive immune response. The IgG molecule has two separate functions; to bind to the |
| | pathogen that elicited the response and to recruit other cells and molecules to destroy |
| | the antigen. The variability of the IgG pool is generated by somatic recombination and |
| | the number of specificities in an individual at a given time point is estimated to be 1011 |
| | variants. |
| | There exists and Nicker |
| | Important Note: |

| This product as supplied is intended for research use only, not for use in human | , |
|--|---|
| therapeutic or diagnostic applications. | |

