

## Rabbit Anti-dog IgG/PE-Cy5.5 antibody

SL0303R-PE-Cy5.5

Chinese Name: PE-Cy5.5标记的兔抗狗IgG  Alias: Immunoglobulin G  Organism Species: Rabbit Clonality: Polyclonal  React Species: dog IF=1:100-1000 Applications: not yet tested in other applications. optimal dilutions/concentrations should be deter  Molecular weight: 150kDa  Form: Lyophilized or Liquid	rmined by the end user.
Organism Species:  Clonality: Polyclonal  React Species:  dog  IF=1:100-1000  Applications: not yet tested in other applications. optimal dilutions/concentrations should be deter  Molecular weight:  150kDa	mined by the end user.
Clonality:  React Species:  dog  IF=1:100-1000  Applications:  not yet tested in other applications.  optimal dilutions/concentrations should be deter  Molecular weight:  150kDa	rmined by the end user.
React Species:    IF=1:100-1000     Applications:   not yet tested in other applications.     optimal dilutions/concentrations should be deter     Molecular weight:   150kDa	mined by the end user.
Applications:  IF=1:100-1000  not yet tested in other applications.  optimal dilutions/concentrations should be deter  Molecular weight:  150kDa	rmined by the end user.
Applications: not yet tested in other applications. optimal dilutions/concentrations should be deter  Molecular weight: 150kDa	rmined by the end user.
optimal dilutions/concentrations should be deter  Molecular weight: 150kDa	rmined by the end user.
Molecular weight: 150kDa	rmined by the end user.
Form: Lyophilized or Liquid	
Concentration: 2mg/1ml	
immunogen: Full length plasma protein:	
Lsotype: IgG	
Purification: affinity purified by Protein A	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Procli	
Storage: Store at –20 oC for one year. Avoid rep	·
Storage: lyophilized antibody is stable at room temperature of the stable at room temperature of the stable at room temperature.	
than a year when kept at -200C. When reconstituted in sterile distilled water or d	
supplied, theantibody is stable for at least two w	
Immunoglobulin G (IgG), is one of the most abu	<u> </u>
levels between 8-17 mg/mL in adult blood. IgG	1
microorganisms and the molecules are produced	
adaptive immune response. The IgG molecule h	
<b>Product Detail:</b> pathogen that elicited the response and to recruit	
the antigen. The variability of the IgG pool is ge	
the number of specificities in an individual at a	given time point is estimated to be 1011
variants.	
Important Note:	

This product as supplied is intended for research use only, not for use in human	,
therapeutic or diagnostic applications.	

