

Rabbit Anti-chicken IgM/APC antibody

SL0314R-APC

| Product Name: | Rabbit Anti-chicken IgM/APC |
|--------------------------|---|
| Chinese Name: | APC标记的兔抗鸡IgM |
| Alias: | Immunoglobulin M |
| Organism Species: | Rabbit |
| Clonality: | Polyclonal |
| React Species: | chk |
| | Flow-Cyt=1:100-1000IF=1:100-1000 |
| Applications: | not yet tested in other applications. |
| | optimal dilutions/concentrations should be determined by the end user. |
| Molecular weight: | 900kDa |
| Form: | Lyophilized or Liquid |
| Concentration: | 2mg/1ml |
| immunogen: | Full length plasma protein: |
| Lsotype: | IgG |
| Purification: | affinity purified by Protein A |
| Storage Buffer: | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage: Product Detail: | Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The |
| | lyophilized antibody is stable at room temperature for at least one month and for greater |
| | than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent |
| | supplied, theantibody is stable for at least two weeks at 2-4 °C. |
| | IgM normally constitutes about 10% of serum immunoglobulins. IgM antibody is |
| | prominent in early immune responses to most antigens and is largely confined to |
| | plasma due to it's large size. Monomeric IgM is expressed as a membrane bound |
| | antibody on the surface of B cells and as a pentamer when secreted by plasma cells. |
| | Due to it's high valency IgM is more efficient than other isotypes is binding antigens |
| | with repeating epitopes (virus particles and red blood cells) and is more efficient than |
| | IgG in activiating the complement pathway. The gene for the mu constant region |
| | contains four domains separated by short intervening sequences. |
| | Tues out out Notes |
| | Important Note: |

| This product as supplied is intended for research use only, not for use in human | , |
|--|---|
| therapeutic or diagnostic applications. | |

