



## Rabbit Anti-FGL2 antibody

SL0315R

<b>Product Name:</b>	FGL2
<b>Chinese Name:</b>	凝血酶原酶/纤维蛋白原2抗体
<b>Alias:</b>	Fibroleukin; Cytotoxic T-lymphocyte-specific protein; Fibrinogen-like protein 2; Fibrinogen like protein 2; Prothrombinase; Fibroleukin; pT49; T49; FGL2_HUMAN.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Rat,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	49kDa
<b>Cellular localization:</b>	Secretory protein
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from Rat FGL2:75-150/422
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	FGL2 is a secreted protein that is similar to the beta- and gamma-chains of fibrinogen. The carboxyl-terminus of the encoded protein consists of the fibrinogen-related domains (FRED). The encoded protein forms a tetrameric complex which is stabilized by interchain disulfide bonds. It may play a role in physiologic functions at mucosal sites. It is constitutively expressed in cytotoxic T-cells. Lack of expression in other lymphoid- and nonlymphoid-derived cell lines suggested that expression of FGL2 may be restricted

to lymphocytes. FGL2 is induced via a mechanism involving IFNG and components of the IFNG signaling pathway, including STAT1 and IRF1.

**Function:**

May play a role in physiologic lymphocyte functions at mucosal sites

**Subunit:**

Homotetramer; disulfide-linked.

**Subcellular Location:**

Secreted.

**Tissue Specificity:**

Constitutively expressed in cytotoxic T-cells.

**Similarity:**

Contains 1 fibrinogen C-terminal domain.

**SWISS:**

N/A

**Gene ID:**

84586

**Database links:**

[Entrez Gene: 511711](#)Cow

[Entrez Gene: 10875](#)Human

[Entrez Gene: 14190](#)Mouse

[Entrez Gene: 84586](#)Rat

[Omim: 605351](#)Human

[SwissProt: Q14314](#)Human

[SwissProt: P12804](#)Mouse

[Unigene: 520989](#)Human


[Unigene: 292100](#)Mouse

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

fgL-2蛋白是新近发现的一种凝血因子, 它通过直接激活凝血酶原启动凝血过程。一

系列实验证明, 该蛋白在肝细胞坏死中起到重要作用。对乙肝患者肝组织免疫组织化学染色hfg12在重型乙型肝炎患者高表达。  
凝血酶原酶(简称fg12)蛋白属于纤维蛋白原家族的一员,由活化的巨噬细胞表达,具有凝血酶原酶的活力,能催化凝血酶原转化为凝血酶,启动凝血过程.



Picture:

Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;  
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;  
Incubation: Anti-fg-12 Polyclonal Antibody, Unconjugated(SL0315R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining