

## Rabbit Anti-Bovine IgM/Alexa Fluor 647 antibody

## SL0327R-AF647

| Product Name:     | Rabbit Anti-Bovine IgM/Alexa Fluor 647  |
|-------------------|---|
| Chinese Name:     | Alexa Fluor 647标记 <b>的兔抗牛</b> IgM   |
| Alias:            | Immunoglobulin M.   |
| Organism Species: | Rabbit  |
| Clonality:        | Polyclonal  |
| React Species:    | cow   |
| Applications:     | Flow-Cyt=1:100-1000IF=1:100-1000  |
|                   | not yet tested in other applications.   |
|                   | optimal dilutions/concentrations should be determined by the end user.                    |
| Molecular weight: | 900kDa  |
| Form:             | Lyophilized or Liquid   |
| Concentration:    | 2mg/1ml   |
| immunogen:        | Full length plasma protein:   |
| Lsotype:          | IgG   |
| Purification:     | affinity purified by Protein A  |
| Storage Buffer:   | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.                          |
| Storage:          | Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The             |
|                   | lyophilized antibody is stable at room temperature for at least one month and for greater |
|                   | than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent  |
|                   | supplied, theantibody is stable for at least two weeks at 2-4 °C.                         |
| Product Detail:   | IgM normally constitutes about 10% of serum immunoglobulins. IgM antibody is              |
|                   | prominent in early immune responses to most antigens and is largely confined to           |
|                   | plasma due to it's large size. Monomeric IgM is expressed as a membrane bound             |
|                   | antibody on the surface of B cells and as a pentamer when secreted by plasma cells.       |
|                   | Due to it's high valency IgM is more efficient than other isotypes is binding antigens    |
|                   | with repeating epitopes (virus particles and red blood cells) and is more efficient than  |
|                   | IgG in activiating the complement pathway. The gene for the mu constant region            |
|                   | contains four domains separated by short intervening sequences.                           |
|                   | Important Note:   |
|                   | important rote.   |

| This product as supplied is intended for research use only, not for use in human | , |
|--|---|
| therapeutic or diagnostic applications.  |   |

