

Goat Anti-Guinea pig IgG/PE-Cy5.5 antibody

SL0358G-PE-Cy5.5

Chinese Name: PE-Cy5.5标记的羊抗豚鼠IgG Alias: Immunoglobulin G Organism Species: Goat Clonality: Polyclonal React Species: Gpig IF=1:100-1000 Applications: optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greated the storage of the storage of the supplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	duct Name:	Goat Anti-Guinea pig IgG/PE-Cy5.5
Organism Species: Clonality: Polyclonal React Species: Gpig IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: I50kDa Form: Lyophilized or Liquid Concentration: immunogen: Lsotype: IgG Purification: Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greateness at year when kept at -20oC. When reconstituted in sterile distilled water or dilusupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	nese Name:	PE-Cy5.5标记 的羊抗豚鼠 IgG
Clonality: Polyclonal React Species: Gpig IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greated than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	as:	Immunoglobulin G
React Species: Gpig IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.	ganism Species:	Goat
Applications: IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: I50kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greated than a year when kept at -20oC. When reconstituted in sterile distilled water or dilusupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	nality:	Polyclonal
Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Lsotype: IgG Purification: Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	ict Species:	Gpig
optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		IF=1:100-1000
Molecular weight:150kDaForm:Lyophilized or LiquidConcentration:2mg/1mlimmunogen:Full length plasma protein:Lsotype:IgGPurification:affinity purified by Protein AStorage Buffer:0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.Storage:Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C.Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	plications:	
Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
Concentration: immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluctions are supplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	lecular weight:	
Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluctional supplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	<u> </u>	
Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for great than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	icentration:	
Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greathen a year when kept at -20oC. When reconstituted in sterile distilled water or diluscular distribution of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	nunogen:	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greather than a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greathan a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
Storage: lyophilized antibody is stable at room temperature for at least one month and for greathan a year when kept at -20oC. When reconstituted in sterile distilled water or diluscupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	rage Buffer:	
than a year when kept at -20oC. When reconstituted in sterile distilled water or dilusupplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
supplied, theantibody is stable for at least two weeks at 2-4 °C. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our	rage:	
Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our		
microorganisms and the molecules are produced by B lymphocytes as a part of our	Product Detail:	
		adaptive immune response. The IgG molecule has two separate functions; to bind to the
		pathogen that elicited the response and to recruit other cells and molecules to destroy
		the antigen. The variability of the IgG pool is generated by somatic recombination and
		the number of specificities in an individual at a given time point is estimated to be 1011
variants.		variants.
Important Note:		Important Note:

This product as supplied is intended for research use only, not for use in human	,
therapeutic or diagnostic applications.	

