



## Goat Anti-Mouse IgM/FITC antibody

SL0368G-FITC

<b>Product Name:</b>	Goat Anti-Mouse IgM/FITC
<b>Chinese Name:</b>	FITC标记的羊抗小鼠IgM
<b>Alias:</b>	Immunoglobulin M.
<b>文献引用</b> <b>PubMed</b> :	<p><b>Specific References(3)</b> SL0368G-FITC has been referenced in 3 publications.</p> <p><b>[IF=3.73]</b>Xin, Jige, et al. "Highly Efficient Generation of GGTA1 Biallelic Knockout Inbred Mini-Pigs with TALENs." PloS one 8.12 (2013): e84250.<b>Pig</b>.  <a href="#">PubMed:24358349</a></p> <p><b>[IF=3.54]</b>Geng, Zhaoxin, et al. "A route to low-cost nanoplasmonic biosensor integrated with optofluidic-portable platform." Sensors and Actuators B: Chemical (2014).<b>other</b>;  <a href="#">PubMed:not posted yet</a></p> <p><b>[IF=4.65]</b>Jiang, P., et al. "Eryptosis as an Underlying Mechanism in Systemic Lupus Erythematosus-Related Anemia." Cellular Physiology and Biochemistry 40.6 (2016): 1391-1400.<b>IF(ICC)</b>;  <a href="#">PubMed:27997909</a></p>
<b>Organism Species:</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	mo
<b>Applications:</b>	Flow-Cyt=1:100-1000IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	900kDa
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	2mg/1ml

<b>immunogen:</b>	Full length plasma protein:
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent supplied, theantibody is stable for at least two weeks at 2-4 °C.
<b>Product Detail:</b>	<p>Immunoglobulin M (IgM) normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent in early immune responses to most antigens and is largely confined to plasma due to it's large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. Due to it's high valency IgM is more efficient than other isotypes is binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activiating the complement pathway. The gene for the mu constant region contains four domains separated by short intervening sequences.</p> <p><b>Important Note:</b> This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p>

www.sunlongbiotech.com