



Rabbit Anti-beta 2 Microglobulin antibody

SL0390R

Product Name:	beta 2 Microglobulin
Chinese Name:	β2-微球蛋白抗体
Alias:	B2M; Beta-2-MG; Beta 2 microglobulin precursor; Beta chain of mhc class 1 proteins; Hdcma22p; B2MG_HUMAN; Beta-2 microglobin; Beta 2 microglobulin precursor; Beta chain of mhc class 1 proteins; Beta chain of MHC class I molecules; Beta-2-microglobulin form pI 5.3; CDABP0092.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,
Applications:	WB=1:500-2000ELISA=1:500-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	13/27kDa
Cellular localization:	Secretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	Recombinant human Beta-2-Microglobulin protein:21-119/119
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	This gene encodes a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has a predominantly beta-pleated sheet structure that can form amyloid fibrils in some pathological conditions. A mutation in this gene has been shown to result in hypercatabolic hypoproteinemia.[provided by RefSeq, Sep 2009].

Function:

Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system.

Subunit:

Heterodimer of an alpha chain and a beta chain. Beta-2-microglobulin is the beta-chain of major histocompatibility complex class I molecules. Polymers of beta 2-microglobulin can be found in tissues from patients on long-term hemodialysis.

Subcellular Location:

Secreted. Note=Detected in serum and urine.

Post-translational modifications:

Glycation of Ile-21 is observed in long-term hemodialysis patients.

DISEASE:

Defects in B2M are the cause of hypercatabolic hypoproteinemia (HYCATHYP) [MIM:241600]. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation.

Note=Beta-2-microglobulin may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. Persistently high beta(2)-microglobulin serum levels lead to amyloidosis in patients on long-term hemodialysis.

Similarity:

Belongs to the beta-2-microglobulin family.

Contains 1 Ig-like C1-type (immunoglobulin-like) domain.

SWISS:

P61769

Gene ID:

567

Database links:

[Entrez Gene: 567](#)Human

[Entrez Gene: 12010](#)Mouse

[Entrez Gene: 24223](#)Rat

[Omim: 109700](#)Human

[SwissProt: P61769](#)Human

[SwissProt: P01887](#)Mouse


[SwissProt: P07151](#)Rat
[Unigene: 534255](#)Human
[Unigene: 702947](#)Human
[Unigene: 163](#)Mouse
[Unigene: 1868](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

小鼠抗人 β 2微球蛋白在中枢神经系统感染、Tumour及全身性自身免疫性疾病患者脑脊液中, 会出现 β 2MG蛋白不同程度的增高。

Picture:



Sample:

Raji(Human) Cell Lysate at 30 ug

Primary: Anti-beta 2 Microglobulin (SL0390R) at 1/2000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 13/27 kD

	Observed band size: 48 kD
--	---------------------------

www.sunlongbiotech.com