

Rabbit anti-MG IgG/Cy7 antibody

SL0403R-Cy7

Chinese Name: Cy7标记的兔抗长爪沙鼠IgG Alias: Immunoglobulin G Organism Species: Rabbit Clonality: Polyclonal React Species: Mongolian Gerbil IF=1:100-1000 Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or disapplied, theantibody is stable for at least two weeks at 2-4 °C.	
Organism Species: Rabbit Clonality: Polyclonal React Species: Mongolian Gerbil IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distille	
Clonality: Polyclonal React Species: Mongolian Gerbil IF=1:100-1000 Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distance.	
React Species: Mongolian Gerbil IF=1:100-1000 Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distance.	
Applications: IF=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distinct the state of the sta	
Applications: not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distinct the storage of the state of the storage of the state	
optimal dilutions/concentrations should be determined by the end user. Molecular weight: 150kDa Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distance.	
Molecular weight:150kDaForm:Lyophilized or LiquidConcentration:2mg/1mlimmunogen:Full length plasma protein:Lsotype:IgGPurification:affinity purified by Protein AStorage Buffer:0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.Storage:Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distinct the constituted in sterile distilled water or distinct the co	
Form: Lyophilized or Liquid Concentration: 2mg/1ml immunogen: Full length plasma protein: Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distil	
Concentration: immunogen: Full length plasma protein: Lsotype: IgG Purification: Storage Buffer: O.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Storag	
immunogen: Lsotype: IgG Purification: Storage Buffer: O.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: S	
Lsotype: IgG Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled	
Purification: affinity purified by Protein A Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or dis	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Storage: Store at -20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled wa	
Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or	
Storage: lyophilized antibody is stable at room temperature for at least one month and for than a year when kept at -20oC. When reconstituted in sterile distilled water or distilled wa	
than a year when kept at -20oC. When reconstituted in sterile distilled water or di	
than a year when kept at -200C. When reconstituted in sterile distilled water or di	
gunnliad thoughthody is stable for at least two weeks at 2.4 °C'	luent
Immunoglobulin G (IgG), is one of the most abundant proteins in serum with nor	
levels between 8-17 mg/mL in adult blood. IgG is important for our defence again	
microorganisms and the molecules are produced by B lymphocytes as a part of our	
adaptive immune response. The IgG molecule has two separate functions; to bind	
Product Detail: pathogen that elicited the response and to recruit other cells and molecules to des	
the antigen. The variability of the IgG pool is generated by somatic recombination	
the number of specificities in an individual at a given time point is estimated to be	3 1011
variants.	
Important Note:	

This product as supplied is intended for research use only, not for use in human	,
therapeutic or diagnostic applications.	

