



SunLong Biotech Co.,LTD
Tel: 0086-571- 56623320 Fax:0086-571- 56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

Rabbit Anti-TIMP-4 antibody

SL0418R

Product Name:	TIMP-4
Chinese Name:	金属蛋白酶组织抑制因子-4抗体
Alias:	Metalloproteinase inhibitor 4; TIMP 4; TIMP4; TIMP metallopeptidase inhibitor 4; TIMP-4; Timp4; TIMP4_HUMAN; Tissue inhibitor of metalloproteinase 4.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Cow,Horse,Rabbit,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	22kDa
Cellular localization:	Secretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human TIMP-4:161-224/224
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	The tissue inhibitors of metalloproteinases (TIMPs) are naturally occurring proteins that specifically inhibit matrix metalloproteinases and regulate extracellular matrix turnover and tissue remodeling by forming tightbinding inhibitory complexes with the MMPs. Thus, TIMPs maintain the balance between matrix destruction and formation. An imbalance between MMPs and the associated TIMPs may play a significant role in the invasive phenotype of malignant tumors. The TIMP proteins share several structural

features including six loops held in place by six disulfide bonds arranged in three knot-like structures. These proteins also contain twelve cysteine residues in conserved regions of the molecule that form six disulfide bonds, essential for the formation of native conformations, and the N terminal region that is necessary for inhibitory activities. The N terminus of each TIMP contains a consensus sequence (VIRAK) and each TIMP is translated with a 29 amino acid leader sequence that is cleaved off to produce the mature protein. The C terminal regions are divergent, which enhances the selectivity of inhibition and binding efficiency. Although the TIMP proteins share high homology, they may either be secreted extracellularly in soluble form (TIMP1, TIMP2 and TIMP4) or bind to extracellular matrix components (TIMP3).

The MMPs and TIMPs can be divided into two groups with respect to gene expression: the majority exhibit inducible expression and a small number are produced constitutively or are expressed at very low levels and are not inducible. Among agents that induce MMP and TIMP production are the inflammatory cytokines TNF alpha and IL1 beta. A marked cell type specificity is a hallmark of both MMP and TIMP gene expression (i.e. a limited number of cell types can be induced to make these proteins). Tissue Inhibitor of Metalloproteinases 4 (TIMP4) was identified by molecular cloning. TIMP4 shows 37 % amino acid identity with TIMP1 and 51 % homology with TIMP2 and TIMP3. TIMP4 is secreted extracellularly, predominantly in heart and brain tissue. It may function in a tissue specific fashion in extracellular matrix (ECM) homeostasis. TIMP4 has a strong inhibitory effect on the invasion of human breast cancer cells across reconstituted basement membranes suggesting that TIMP4 may have an important role in inhibiting primary tumor growth and progression. The human TIMP4 gene has the chromosomal location of 3p25.

Function:

Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7 and MMP-9.

Subcellular Location:

Secreted.

Tissue Specificity:

Abundant in heart and present at low levels in many other tissues.

Similarity:

Belongs to the protease inhibitor I35 (TIMP) family.

Contains 1 NTR domain.

SWISS:

Q99727

Gene ID:

7079

Database links:

[Entrez Gene: 7079](#)Human

[Entrez Gene: 110595](#)Mouse

[Entrez Gene: 114849](#)Rat

[Entrez Gene: 680130](#)Rat

[Omim: 601915](#)Human

[SwissProt: Q99727](#)Human

[SwissProt: Q9JHB3](#)Mouse

[SwissProt: P81556](#)Rat

[Unigene: 591665](#)Human

[Unigene: 255607](#)Mouse

[Unigene: 155651](#)Rat


Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


Synthesis and Degradation (Synthesis and Degradation)

TIMP-4是TIMP家族的新成员, TIMP-3与TIMP-

4同样具有较强的促凋亡、抑制血管生成、抗Tumour浸润作用。

Picture:

Paraformaldehyde-fixed, paraffin embedded (Rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TIMP-4) Polyclonal Antibody, Unconjugated (SL0418R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: human endometrium carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-TIMP-4 Polyclonal Antibody, Unconjugated(SL0418R) 1:300, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining

www.sunlongbiotech.com