



Rabbit Anti-Chicken IgY/HRP antibody

SL0432R-HRP

Product Name:	Rabbit Anti-Chicken IgY/HRP
Chinese Name:	辣根过氧化物酶标记兔抗鸡IgY抗体
文献引用 PubMed :	<p>Specific References(1) SL0432R-HRP has been referenced in 1 publications.</p> <p>[IF=1.73]Liu, Dong, and Zhong-Xiang Niu. "Cloning of a gene fragment encoding chicken complement component C3d with expression and immunogenicity of Newcastle disease virus F gene?CC3d fusion protein." Avian Pathology 37.5 (2008): 477-485. iWB;Chicken.</p> <p style="text-align: right;">PubMed:8798021</p>
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Chicken,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	Chicken IgY:
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Product Detail:	In chickens, immunoglobulin Y is the functional equivalent to Immunoglobulin G (IgG). Like IgG, it is composed of two light and two heavy chains. Structurally, these two types of immunoglobulin differ primarily in the heavy chains, which in IgY have a

molecular mass of about 65,100 atomic mass units (amu), and are thus larger than in IgG. The light chains in IgY, with a molar mass of about 18,700 amu, are somewhat smaller than the light chains in IgG. The molar mass of IgY thus amounts to about 167,000 amu. The steric flexibility of the IgY molecule is less than that of IgG. Functionally, IgY is partially comparable to Immunoglobulin E (IgE), as well as to IgG. However, in contrast to IgG, IgY does not bind to Protein A, to Protein G, or to cellular Fc receptors. Furthermore, IgY does not activate the complement system. The name Immunoglobulin Y was suggested in 1969 by G.A. Leslie and L.W. Clem, after they were able to show differences between the immunoglobulins found in chicken eggs, and immunoglobulin G. Other synonymous names are Chicken IgG, Egg Yolk IgG, and 7S-IgG.

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com