



## Rabbit Anti-VSIG4 antibody

SL0479R

<b>Product Name:</b>	VSIG4
<b>Chinese Name:</b>	Tlymphocyte负调节蛋白抗体
<b>Alias:</b>	V-set and immunoglobulin domain-containing protein 4; CRIG; V set and immunoglobulin domain containing 4; Z39IG; VSIG4_HUMAN.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Dog,Cow,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	33kDa
<b>Cellular localization:</b>	cytoplasmicThe cell membrane
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human VSIG4:81-160/399
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	T cell activation by APCs is positively and negatively regulated by members of the B7 family. We have identified a previously unknown function for B7 familyrelated protein V-set and Ig domaincontaining 4 (VSIG4). Administration to mice of soluble VSIG4-Ig fusion molecules reduced the induction of T cell responses in vivo and inhibited the production of Th celldependent IgG responses. Unlike that of B7 family members, surface expression of VSIG4 was restricted to resting tissue macrophages and absent

upon activation by LPS or in autoimmune inflammatory foci. The specific expression of VSIG4 on resting macrophages in tissue suggests that this inhibitory ligand may be important for the maintenance of T cell unresponsiveness in healthy tissues.

**Function:**

Phagocytic receptor, strong negative regulator of T-cell proliferation and IL2 production. Potent inhibitor of the alternative complement pathway convertases.

**Subcellular Location:**

Membrane; Single-pass type I membrane protein (Probable).

**Tissue Specificity:**

Abundantly expressed in several fetal tissues. In adult tissues, highest expression in lung and placenta. Expressed in resting macrophages.

**Similarity:**

Contains 2 Ig-like (immunoglobulin-like) domains.

**SWISS:**

Q9Y279

**Gene ID:**

11326

**Database links:**

[Entrez Gene: 11326](#)Human

[Entrez Gene: 278180](#)Mouse

[Entrez Gene: 312102](#)Rat

[Omim: 300353](#)Human

[SwissProt: Q9Y279](#)Human

[Unigene: 8904](#)Human


[Unigene: 26781](#)Mouse

[Unigene: 138093](#)Rat

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Sample:


Lung (Mouse) Lysate at 40 ug

Primary: Anti-VSIG4 (SL0479R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 33 kD

Observed band size: 33 kD



Sample:

Lung(Mouse)Lysate at 30 ug


Placenta(Mouse) Lysate at 30 ug

Primary: Anti-VSIG4 (SL0479R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 33 kD

Observed band size: 33 kD



Sample:


Placenta (Mouse) Lysate at 40 ug

Primary: Anti-VSIG4 (SL0479R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 33 kD

Observed band size: 33 kD



Tissue/cell: Rat lung tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;  
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block  
endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer  
(normal goat serum,C-0005) at 37°C for 20 min;  
Incubation: Anti- VSIG4 Polyclonal Antibody, Unconjugated(SL0479R) 1:200,  
overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and  
DAB(C-0010) staining