



## Rabbit Anti-FSH receptor antibody

SL0895R

<b>Product Name:</b>	FSH receptor
<b>Chinese Name:</b>	促卵泡刺激素受体抗体
<b>Alias:</b>	Follicle-stimulating hormone receptor precursor; Follicle stimulating hormone receptor; Follicle stimulating hormone receptor isoform 1; Follitropin receptor; FSHRO; LGR1; ODG1; Follicle-stimulating hormone receptor; FSH-R; FSH R; Fshr; FSHR_HUMAN; MGC141667; MGC141668; ovarian dysgenesis 1.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Dog,Pig,Cow,Horse,Sheep,Guinea Pig,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800Flow-Cyt=1µg/TestIF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	78kDa
<b>Cellular localization:</b>	The cell membrane
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human FSHR:201-300/695<Extracellular>
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	The protein encoded by this gene belongs to family 1 of G-protein coupled receptors. It is the receptor for follicle stimulating hormone and functions in gonad development. Mutations in this gene cause ovarian dysgenesis type 1, and also ovarian

hyperstimulation syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2010]

**Function:**

Receptor for follicle-stimulating hormone. The activity of this receptor is mediated by G proteins which activate adenylate cyclase.

**Subunit:**

Interacts with ARRB2.

**Subcellular Location:**

Cell membrane; Multi-pass membrane protein.

**Tissue Specificity:**

Sertoli cells and ovarian granulosa cells.

**DISEASE:**

Defects in FSHR are a cause of ovarian dysgenesis type 1 (ODG1) [MIM:233300]; also known as premature ovarian failure or gonadal dysgenesis XX type or XX gonadal dysgenesis (XXGD) or hereditary hypergonadotropic ovarian failure or hypergonadotropic ovarian dysgenesis with normal karyotype. ODG1 is an autosomal recessive disease characterized by primary amenorrhea, variable development of secondary sex characteristics, and high serum levels of follicle-stimulating hormone (FSH) and luteinizing hormone (LH).

Defects in FSHR are a cause of ovarian hyperstimulation syndrome (OHSS) [MIM:608115]. OHSS is a disorder which occurs either spontaneously or most often as an iatrogenic complication of ovarian stimulation treatments for in vitro fertilization. The clinical manifestations vary from abdominal distention and discomfort to potentially life-threatening, massive ovarian enlargement and capillary leak with fluid sequestration. Pathologic features of this syndrome include the presence of multiple serous and hemorrhagic follicular cysts lined by luteinized cells, a condition called hyperreactio luteinalis.

**Similarity:**

Belongs to the G-protein coupled receptor 1 family. FSH/LSH/TSH subfamily. Contains 9 LRR (leucine-rich) repeats. Contains 1 LRRNT domain.

**SWISS:**

P23945

**Gene ID:**

2492

**Database links:**

[Entrez Gene: 281172](#)Cow

[Entrez Gene: 100861291](#)Goat

[Entrez Gene: 2492](#)Human

[Entrez Gene: 14309](#)Mouse

[Entrez Gene: 397679](#)Pig

[Entrez Gene: 25449](#)Rat

[Entrez Gene: 443299](#)Sheep

[Omim: 136435](#)Human

[SwissProt: P35376](#)Cow

[SwissProt: P23945](#)Human

[SwissProt: P35378](#)Mouse

[SwissProt: P49059](#)Pig

[SwissProt: P20395](#)Rat

[SwissProt: P35379](#)Sheep

[Unigene: 1428](#)Human

[Unigene: 57155](#)Mouse


[Unigene: 162843](#)Rat

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

FSHR属于G protein-coupled receptor超家族中的glycoprotein亚家族成员, Transmembrane protein。人类的FSHR是由695个氨基酸组成的一条肽链。主要分布于睾丸支持细胞和卵巢颗粒细胞, 成熟的FSHR蛋白分子质量为:75kDa。

Picture:



Sample:


Testis (Mouse) Lysate at 40 ug

Primary: Anti-FSH receptor (SL0895R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 78 kD

Observed band size: 78 kD



**bs-0895R Anti-FSH receptor Polyclonal Antibody**


Tissue/cell: rat ovary tissue; 4% Paraformaldehyde-fixed and paraffin-embedded

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min

Block endogenous peroxidase by 3% Hydrogen peroxide for 30min


Blocking buffer (normal goat serum) at 37°C for 20 min

Incubation: Anti-FSH receptor Polyclonal Antibody, Unconjugated(bs-0895R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining



Paraformaldehyde-fixed, paraffin embedded (rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for


30min; Antibody incubation with (FSH receptor) Polyclonal Antibody, Unconjugated (SL0895R) at 1:200 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: human placenta tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-FSH receptor Polyclonal Antibody, Unconjugated(SL0895R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: HUVEC cells(blue).

Primary Antibody:Rabbit Anti-FSH receptor antibody(SL0895R), Dilution: 1 $\mu$ g in 100  $\mu$ L 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions );

Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

#### Protocol

The cells were fixed with 2% paraformaldehyde (10 min) .Primary antibody (SL0895R) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 1 0% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Anti-rabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.