



Rabbit Anti-Leptin receptor antibody

SL0961R

Product Name:	Leptin receptor
Chinese Name:	瘦素受体抗体
Alias:	obl; CD 295; CD295; CD295 antigen; Db; Fa; HuB219; LEP R; LEPR; LEPROT; Leptin receptor fatty; Leptin receptor gene related protein; Leptin receptor precursor; OB R; OB-R; OB R gene related protein; OB receptor; OB-RGRP; Obr; Ob-R a/b/c/d/e; LEPR HUMAN; LEPRD; Leptin receptor.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep,
Applications:	WB=1:500-2000IHC-P=1:400-800IHC-F=1:400-800Flow-Cyt=1 μ g/TestIF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	130kDa
Cellular localization:	The cell membraneSecretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human Leptin receptor:501-600/1165<Extracellular>
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	The protein encoded by this gene belongs to the gp130 family of cytokine receptors that are known to stimulate gene transcription via activation of cytosolic STAT proteins. This protein is a receptor for leptin (an adipocyte-specific hormone that regulates body

weight), and is involved in the regulation of fat metabolism, as well as in a novel hematopoietic pathway that is required for normal lymphopoiesis. Mutations in this gene have been associated with obesity and pituitary dysfunction. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. It is noteworthy that this gene and LEPROT gene (GeneID:54741) share the same promoter and the first 2 exons, however, encode distinct proteins (PMID:9207021).[provided by RefSeq, Nov 2010]

Function:

Receptor for obesity factor (leptin). On ligand binding, mediates signaling through JAK2/STAT3. Involved in the regulation of fat metabolism and, in a hematopoietic pathway, required for normal lymphopoiesis. May play a role in reproduction. Can also mediate the ERK/FOS signaling pathway.

Subunit:

On leptin stimulation, homodimerizes. The phosphorylated receptor binds a number of SH2 domain-containing proteins such as STAT3, PTPN11, and SOCS3. Interaction with SOCS3 inhibits LRB signaling.

Subcellular Location:

Cell membrane; Single-pass type I membrane protein. Isoform E: Secreted (Probable).

Tissue Specificity:

Isoform A is expressed in fetal liver and in hematopoietic tissues and choroid plexus. In adults highest expression in heart, liver, small intestine, prostate and ovary. Low level in lung and kidney. Isoform B is highly expressed in hypothalamus.

Post-translational modifications:

On ligand binding, phosphorylated on two conserved C-terminal tyrosine residues (isoform B only) by JAK2. Tyr-986 is required for complete binding and activation of PTPN11, ERK/FOS activation and, for interaction with SOCS3. Phosphorylation on Tyr-1141 is required for STAT3 binding/activation.

Similarity:

Belongs to the type I cytokine receptor family. Type 2 subfamily.
Contains 4 fibronectin type-III domains.
Contains 1 Ig-like (immunoglobulin-like) domain.

SWISS:

P48357

Gene ID:

3953

Database links:

[Entrez Gene: 3953](#) Human

[Entrez Gene: 16847](#) Mouse

[Entrez Gene: 24536](#) Rat

[Omim: 601007](#) Human

[SwissProt: P48357](#) Human

[SwissProt: P48356](#) Mouse

[SwissProt: Q62959](#) Rat

[Unigene: 23581](#) Human

[Unigene: 723178](#) Human

[Unigene: 259282](#) Mouse


[Unigene: 9891](#) Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Leptin是一种分子量为16kDa的脂肪组织源激素,由脂肪细胞分泌的,具有强亲水性,以单体形式存在于血浆中。Leptin具有广泛的生物学效应,作用于下丘脑,调节食欲、能量代谢及体重。Leptin还可能作为脂肪-胰岛内分泌轴一部分,参与胰岛素分泌的调节。Leptin receptor (OB-R)是与gp130最密切相关的Signal transduction的共同成分,属1类cell factor超家族,下丘脑有丰富的Leptin receptor,在其它部位也有分布,但水平较低,如:胰岛素B细胞、脉络丛、肝、肺、心、肾、睾丸、lymphocyte和脂肪细胞中。人类存在四个异性体,均为单跨膜受体。其中OB-Rb在胞内结构域最长,是唯一能进行信号传导并调节热量摄入和能量消耗的异性体,分子量为132kDa。

Picture:



Sample:


Liver(Mouse) Lysate at 40 ug

Primary: Anti-Leptin receptor? (SL0961R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 130 kD

Observed band size: 155 kD



Sample:

Liver (Rat) Lysate at 40 ug


Primary: Anti-Leptin receptor? (SL0961R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution


Predicted band size: 130 kD

Observed band size: 160 kD

www.sunlongbiotech.com



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LEPR) Polyclonal Antibody, Unconjugated (SL0961R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Tissue/cell: mouse spleen tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Leptin receptor Polyclonal Antibody, Unconjugated(SL0961R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-

0023) and DAB(C-0010) staining



Primary Antibody: Rabbit Anti-Leptin receptor antibody(SL0961R)

Isotype Control Antibody: Rabbit IgG(orange) Secondary Antibody: F(ab')₂ fragment goat anti-rabbit IgG-PE(white blue)

Positive control: K562 cells(Blue)(An equivalent amount of pre-warmed 4% paraformaldehyde was added and the cells were incubated for 10 min at 37 °C.)