



Rabbit Anti-Caspase-12 antibody

SL0967R


Product Name:	Caspase-12
Chinese Name:	半胱氨酸蛋白酶蛋白-12抗体
Alias:	Caspase-12; UNQ9415; Apoptosis related cysteine protease; CASP 12; casp12; CASP12P1; Caspase 12 pseudogene 1; CASP-12; casp12; caspase 12 (gene/pseudogene); Caspase12; CASPC_HUMAN; OTTHUMP00000207026; OTTHUMP00000207028; OTTHUMP00000207031; OTTHUMP00000207032; OTTHUMP00000219058; UNQ9415.
文献引用 PubMed :	Specific References(1) SL0967R has been referenced in 1 publications. [IF=3.53] Fang C, Zhang J, Qi D, Fan X, Luo J, et al. (2014) Evodiamine Induces G2/M Arrest and Apoptosis via Mitochondrial and Endoplasmic Reticulum Pathways in H446 and H1688 Human Small-Cell Lung Cancer Cells. PLoS ONE 9(12): e115204. WB;Rat. PubMed:25506932
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	38kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from the middle of human Caspase-12:161-250/341
Lsotype:	IgG
Purification:	affinity purified by Protein A

Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their substrate molecules. This gene is most highly related to members of the ICE subfamily of caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates apoptosis in response to endoplasmic reticulum stress. However, in humans this gene contains a polymorphism for the presence or absence of a premature stop codon. The majority of human individuals have the premature stop codon and produce a truncated non-functional protein. The read-through codon occurs primarily in individuals of African descent and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe sepsis. Several alternatively spliced transcript variants have been noted for this gene. [provided by RefSeq, Feb 2011]</p> <p>Function: Has no protease activity. May reduce cytokine release in response to bacterial lipopolysaccharide during infections. Reduces activation of NF-kappa-B in response to TNF.</p> <p>Tissue Specificity: Detected in heart, kidney, liver, lung, pancreas, small intestine, spleen, stomach, thymus and testis.</p> <p>Similarity: Belongs to the peptidase C14A family. Contains 1 CARD domain.</p> <p>SWISS: Q6UXS9</p> <p>Gene ID: 100506742</p> <p>Database links: Entrez Gene: 100506742 Human SwissProt: Q6UXS9 Human Unigene: 476989 Human</p> <p>Important Note:</p>

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

ASPASE-

12是caspase家族成员,是存在于内质网上的促凋亡因子,内质网应激早期可通过GRP(glucose-regulated protein, GRP)的表达增多来保护细胞;CASPASE-12是内质网细胞关键介导因子, caspase-12还是caspase-1的抑制剂。



Picture:

Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Caspase-12 Polyclonal Antibody, Unconjugated(SL0967R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining