




## Rabbit Anti-NQO2 antibody

SL10584R

<b>Product Name:</b>	NQO2
<b>Chinese Name:</b>	NAD(P)H 苯醌脱氢酶 2抗体
<b>Alias:</b>	DHQV; DIA6; EC 1.10.99.2; MGC94180; NAD(P)H dehydrogenase quinone 2; NAD(P)H menadione oxidoreductase 1 dioxin inducible 2; NAD(P)H menadione oxidoreductase 2 dioxin inducible; NMOR2; NQO 2; NQO2; NQO2_HUMAN; NRH dehydrogenase [quinone] 2; NRH dehydrogenase; NRH:quinone oxidoreductase 2; OTTHUMP00000015948; OTTHUMP00000015949; OTTHUMP00000015953; Ox 2; Ox2; QR2; Quinone; Quinone reductase 2; Ribosyldihydronicotinamide dehydrogenase [quinone]; Ribosyldihydronicotinamide dehydrogenase.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Dog,Rabbit,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	26kDa
<b>Cellular localization:</b>	cytoplasmic
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human NQO2:11-110/231
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	

Picture:



Sample:

liver (mouse) Lysate at 40 ug

brain (Mouse) Lysate at 40 ug

Primary: Anti-NQO2(SL10584R) at 1/300 dilution

Secondary: HRP conjugated Goat-Anti-rabbit IgG(SL10584R) at 1: 5000;

Predicted band size: 26 kD

Observed band size: 26 kD