



Rabbit Anti-Myeloperoxidase antibody

SL1061R

Product Name:	Myeloperoxidase
Chinese Name:	髓过氧化物酶抗体
Alias:	Myeloperoxidase; MPO; c-ANCA; 89 kDa myeloperoxidase; 84 kDa yeloperoxidase; Myeloperoxidase light chain; Myeloperoxidase heavy chain; EC 1.11.1.7; PERM HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Cow,Horse,Rabbit,Guinea Pig,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800Flow-Cyt=1ug/TestIF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	84kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human Myeloperoxidase:51-150/745
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	Myeloperoxidase (MPO) is a heme protein synthesized during myeloid differentiation that constitutes the major component of neutrophil azurophilic granules. Produced as a single chain precursor, myeloperoxidase is subsequently cleaved into a light and heavy chain. The mature myeloperoxidase is a tetramer composed of 2 light chains and 2 heavy chains. This enzyme produces hypohalous acids central to the microbicidal activity of

netrophils. [provided by RefSeq, Jul 2008].

Function:

Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity.

Subunit:

Tetramer of two light chains and two heavy chains.

Subcellular Location:

Lysosome.

DISEASE:

Defects in MPO are the cause of myeloperoxidase deficiency (MPD) [MIM:254600]. MPD is an autosomal recessive defect that results in disseminated candidiasis.

Similarity:

Belongs to the peroxidase family, XPO subfamily.

SWISS:

P05164

Gene ID:

4353

Database links:

[Entrez Gene: 4353](#) Human

[Entrez Gene: 17523](#) Mouse

[Entrez Gene: 303413](#) Rat

[Omim: 606989](#) Human

[SwissProt: P05164](#) Human

[SwissProt: P11247](#) Mouse

[Unigene: 458272](#) Human

[Unigene: 4668](#) Mouse

[Unigene: 47782](#) Rat


Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

髓过氧化物酶MPO,作为一种白细胞酶,具有介导炎性反应、调节免疫应答等多种功能,并可参与疾病的发生发展过程。同时,髓过氧化物酶基因存在基因多态性,也影响机体对疾病的易感性。

在正常淋巴组织中和各种髓样细胞增生症中, MPO均有较强表达, 如:淋巴样细胞、原核细胞、肥大细胞、浆细胞以及各种上皮源性Tumour和肉瘤等。

Picture:



Sample:


Liver (Mouse) Lysate at 40 ug

Primary: Anti- Myeloperoxidase (SL1061R) at 1/300 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 84 kD

Observed band size: 63 kD




Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Myeloperoxidase) Polyclonal Antibody, Unconjugated (SL1061R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;


Incubation: Anti-Myeloperoxidase Polyclonal Antibody, Unconjugated(SL1061R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse lymphoma tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-MPO/c-ANCA Polyclonal Antibody, Unconjugated(SL1061R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: HL60.

Primary Antibody (green line): Rabbit Anti-Myeloperoxidase antibody (SL1061R)

Dilution: 1 μ g /10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 μ g /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature.

Acquisition of 20,000 events was performed.