

# Rabbit Anti-COLEC12 antibody

## SL10736R

Product Name:	COLEC12
Chinese Name:	凝集胎盘蛋白1抗体
Alias:	Collectin placenta protein 1; Collectin placenta 1; Collectin-12; hCL-P1; CL-P1; Nurse cell scavenger receptor; Scavenger receptor with C-type lectin; Scavenger receptor class A member 4; COLEC12; CLP1; NSR2; SCARA4; SRCL. COL12 HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human, Mouse, Rat, Dog, Pig, Cow, Rabbit,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	81kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human COLEC12:31-130/742 <extracellular></extracellular>
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	<u>PubMed</u>
Product Detail:	Colec12 is a member of the C-lectin family, proteins that possess collagen-like sequences and carbohydrate recognition domains. Colec12 is a scavenger receptor that displays several functions associated with host defense. It promotes binding and phagocytosis of Gram-positive, Gram-negative bacteria and yeast and mediates the

recognition, internalization and degradation of oxidatively modified low density lipoprotein (oxLDL) by vascular endothelial cells.

#### Function:

Scavenger receptor that displays several functions associated with host defense. Promotes binding and phagocytosis of Gram-positive, Gram-negative bacteria and yeast. Mediates the recognition, internalization and degradation of oxidatively modified low density lipoprotein (oxLDL) by vascular endothelial cells. Binds to several carbohydrates including Gal-type ligands, D-galactose, L- and D-fucose, GalNAc, T and Tn antigens in a calcium-dependent manner and internalizes specifically GalNAc in nurse-like cells. Binds also to sialyl Lewis X or a trisaccharide and asialo-orosomucoid (ASOR). May also play a role in the clearance of amyloid beta in Alzheimer disease.

#### **Subunit:**

The extracellular domain forms a stable trimer. The extracellular domain interacts with fibrillar beta amyloid peptide.

#### **Subcellular Location:**

Membrane; Single-pass type II membrane protein. Note=Forms clusters on the cell surface.

#### Tissue Specificity:

Expressed in perivascular macrophages. Expressed in plaques-surrounding reactive astrocytes and in perivascular astrocytes associated with cerebral amyloid angiopathy (CAA) in the temporal cortex of Alzheimer patient (at protein level). Strongly expressed in placenta. Moderately expressed in heart, skeletal muscle, small intestine and lung. Weakly expressed in brain, colon, thymus and kidney. Expressed in nurse-like cells. Expressed in reactive astrocytes and vascular/perivascular cells in the brain of Alzheimer patient.

#### Similarity:

Contains 1 C-type lectin domain. Contains 3 collagen-like domains.

#### **SWISS:**

Q5KU26

#### Gene ID:

81035

#### Database links:

Entrez Gene: 421061 Chicken

Entrez Gene: 504741 Cow

Entrez Gene: 81035Human

Entrez Gene: 140792Mouse

Entrez Gene: 361289Rat

Omim: 607621Human

SwissProt: Q2LK54Chicken

SwissProt: A6QP79Cow

SwissProt: Q5KU26Human

SwissProt: Q8K4Q8Mouse

SwissProt: Q4V885Rat

<u>Unigene: 33741</u>Chicken

Unigene: 4492Cow

Unigene: 464422Human

Unigene: 218571Mouse

Unigene: 31273Rat

### Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.