



## Rabbit Anti-PCDHGA7 antibody

SL11148R

<b>Product Name:</b>	PCDHGA7
<b>Chinese Name:</b>	原钙粘蛋白 $\gamma$ A7抗体
<b>Alias:</b>	PCDG7_HUMAN; PCDH-gamma-A7; PCDHGA7; protocadherin gamma subfamily A, 7; Protocadherin gamma-A7.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,
<b>Applications:</b>	IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	99kDa
<b>Cellular localization:</b>	The cell membrane
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human PCDHGA7:301-400/932
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin (PCDH) gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDHGA7 (Protocadherin gamma-A7) is a 932 amino acid that is one of 22 proteins encoded by the protocadherin gamma cluster. The protocadherein gamma cluster consists of three

subfamilies (A, B and C) and PCDHGA7 is a member of the gamma subfamily A. PCDHGA7 is a type I transmembrane receptor containing six cadherin motifs and is expressed in the central nervous system where it localizes to synapses. Members of the gamma cluster of protocadherins are essential for neuronal survival. There are two isoforms of PCDHGA7 that are produced as a result of alternative splicing events.

**Subcellular Location:**

Cell membrane; Single-pass type I membrane protein

**Similarity:**

Contains 6 cadherin domains.

**SWISS:**

Q9Y5G6

**Gene ID:**

56108

**Database links:**

[Entrez Gene: 56108](#)Human

[SwissProt: Q9Y5G6](#)Human

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.