



Rabbit Anti-PCDHGB2 antibody

SL11156R

Product Name:	PCDHGB2
Chinese Name:	原钙粘蛋白 γ B2抗体
Alias:	PCDH-gamma-B2; Protocadherin gamma B2; PCDGE HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Pig,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	98kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human PCDHGB2:21-120/931<Extracellular>
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. The protocadherein gamma cluster consists of three subfamilies (A, B and C). As a member of the gamma subfamily B, PCDHGB2 (Protocadherin gamma B2) is a 931 amino acid protein that is one of 22

proteins encoded by the protocadherin gamma cluster. Typical of gamma protocadherins, PCDHGB2 contains six cadherin motifs and is a type I transmembrane receptor expressed in the central nervous system. With localization to synapses, members of the gamma cluster of protocadherins are essential for neuronal survival. There are two isoforms of PCDHGB2 that are produced as a result of alternative splicing events.

Function:

Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Subcellular Location:

Cell Membrane

Similarity:

Contains 6 cadherin domains.

SWISS:

Q9Y5G2

Gene ID:

56103

Database links:

[Entrez Gene: 56103](#)Human

UniProtKB/Swiss-Prot: Q9Y5G2.1

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.