



## Rabbit Anti-AMIGO2 antibody

SL11450R

<b>Product Name:</b>	AMIGO2
<b>Chinese Name:</b>	粘附分子IgG样结构域蛋白2抗体
<b>Alias:</b>	Adhesion molecule with Ig like domain 2; ALI 1; ALI1; Alivin 1; Alivin-1; Alivin1; AMIGO 2; Amphoterin induced gene 2; Amphoterin induced protein 2; DEGA; AMGO2_HUMAN; AMIGO-2; Amigo2; Amphoterin induced protein 2; Amphoterin-induced protein 2; Differentially expressed in gastric adenocarcinoma; Differentially expressed in gastric adenocarcinomas; Transmembrane protein AMIGO 2; Transmembrane protein AMIGO2.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Dog,Pig,Cow,Horse,Sheep,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	54kDa
<b>Cellular localization:</b>	The nucleusThe cell membrane
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human AMIGO2:21-120/522<Extracellular>
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	The amphoterin-induced gene and ORF (AMIGO) family of proteins consists of

AMIGO-1, AMIGO-2 and AMIGO-3. All three members are single pass type I membrane proteins that contain several leucine-rich repeats, one IgG domain, and a transmembrane domain. The AMIGO proteins are specifically expressed on fiber tracts of neuronal tissues and participate in their formation. The AMIGO proteins can form complexes with each other, but can also bind itself. AMIGO-1, also designated Alivin-2, promotes growth and fasciculation of neurites and plays a role in myelination and fasciculation of developing neural axons. In cerebellar neurons, AMIGO-2 (Alivin-1) is crucial for depolarization-dependent survival. Similar to AMIGO-1 and AMIGO-2, AMIGO-3 (Alivin-3) plays a role in homophilic and/or heterophilic cell-cell interaction and signal transduction.

**Function:**

A novel primary response gene, AMIGO2, is an activity-dependent gene and promotes survival of neurons. Sequence analyses reveal that rat, mouse, and human AMIGO2 proteins contain seven leucine-rich repeats, one IgC2-like loop and a transmembrane domain, and display homology to Kek and Trk families. Results suggest that expression of ali1 promotes depolarization-dependent survival of the granule neuron. Mouse AMIGO2 was mapped to a locus approximately 55.3 cm from the centromere on chromosome 15 that is syntenic to positional candidate loci for familial Alzheimer's disease type 5 and Parkinson's disease 8 on human chromosome 12.

**Subunit:**

Binds itself as well as AMIGO1 and AMIGO3 (By similarity).

**Subcellular Location:**

Cell membrane; Single-pass type I membrane protein. Nucleus. Note: Associated with nucleus as well as plasma membrane. Restricted to somata of cerebellar as well as hippocampal neurons By

**Tissue Specificity:**

Highest levels in breast, ovary, cervix, and uterus. Lower levels in lung, colon, and rectum. Differentially expressed in 56% of thyroid, 57% of pancreatic and 45% of stomach cancers.

**Similarity:**

Belongs to the immunoglobulin superfamily. AMIGO family.  
Contains 1 Ig-like C2-type (immunoglobulin-like) domain.  
Contains 6 LRR (leucine-rich) repeats.  
Contains 1 LRRCT domain.  
Contains 1 LRRNT domain.

**SWISS:**

Q86SJ2

**Gene ID:**

347902

**Database links:**

[Entrez Gene: 347902](#) Human


[SwissProt: Q86SJ2](#) Human

[Unigene: 121520](#) Human

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Picture:**



Sample:


Hela(Human) Cell Lysate at 30 ug

Primary: Anti- AMIGO2 (SL11450R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 54 kD

Observed band size: 54 kD




Tissue/cell: mouse placenta tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-AMIGO2 Polyclonal Antibody, Unconjugated(SL11450R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (human breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AMIGO2) Polyclonal Antibody, Unconjugated (SL11450R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.