



Rabbit Anti-TAGLN3 antibody

SL12041R

Product Name:	TAGLN3
Chinese Name:	神经元蛋白22抗体
Alias:	Neuronal protein 22; Neuronal protein NP25; NP22; NP25; TAGL3_HUMAN; TAGLN3; Transgelin 3; Transgelin-3.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Chicken,Dog,Pig,Cow,Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	22kDa
Cellular localization:	The nucleus
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human TAGLN3/NP22:8-110/199
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	Transgelin is a homolog of transgelin and is also expressed in smooth muscle cells and by peritoneal B-1 cells. Unlike the other two transgelin proteins, transgelin-3 (also designated TAGLN2, NP22 (neuronal protein 22) or NP25) is predominantly expressed in brain. Transgelin-3 contains a putative Actin-binding domain, two EF-hand motifs, two potential phosphorylation sites and a calponin-homology (CH) domain. Transgelin-3 shares homology with transgelin and Calponin, two cytoskeleton-interacting proteins.

Belonging to the calponin family, transgelin-3 colocalizes with Actin and Tubulin, suggesting a possible role for transgelin-3 in neuronal plasticity or as a signaling protein. Due to a varied expression pattern, transgelin-3 may play different roles in the developing and adult brain. Expression of transgelin-3 is upregulated in regions of the human alcoholic brain.

Tissue Specificity:

Widely expressed in the brain. Expression is increased in the superior frontal cortex of alcoholics, but not in the motor cortex or cerebellum.

Similarity:

Belongs to the calponin family.
Contains 1 calponin-like repeat.
Contains 1 CH (calponin-homology) domain.

SWISS:

Q9UI15

Gene ID:

29114

Database links:

[Entrez Gene: 29114](#) Human

[Entrez Gene: 56370](#) Mouse

[Entrez Gene: 63837](#) Rat

[Entrez Gene: 515562](#) Cow

[Omim: 607953](#) Human

[SwissProt: Q3ZBY2](#) Cow

[SwissProt: Q9UI15](#) Human

[SwissProt: Q9R1Q8](#) Mouse

[SwissProt: P37805](#) Rat


[Unigene: 169330](#) Human

[Unigene: 24183](#) Mouse

[Unigene: 10998](#) Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-TAGLN3 Polyclonal Antibody, Unconjugated(SL12041R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining