

Rabbit Anti-ZBED2 antibody

SL13552R

einBED抗体
BED2; ZBED2_HUMAN; Zinc finger BED domain containing protein ED domain-containing protein 2; Zinc finger, BED domain containing
eep,
0ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-00 (Paraffin sections need antigen repair) other applications. s/concentrations should be determined by the end user.
·
Liquid
d synthetic peptide derived from human ZBED2:1-100/218
by Protein A
7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
for one year. Avoid repeated freeze/thaw cycles. The lyophilized le at room temperature for at least one month and for greater than a year 0°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of libody is stable for at least two weeks at 2-4 °C.
xin-binding protein involved in Wnt/beta-catenin signaling modulation. ion has been shown in human and mouse retinas where it is thought to bry protein in cone photoreceptors and Müller cells. The functions of ED5 have yet to be elucidated.
)

Similarity:

Contains 1 BED-type zinc finger.

SWISS: Q9BTP6

Gene ID: 79413

Database links:

Entrez Gene: 79413 Human

SwissProt: Q9BTP6 Human

Unigene: 136912 Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.