

Rabbit Anti-DDX56 antibody

SL14237R

Product Name:	DDX56	
Chinese Name:	ATP依赖RNA解旋酶DDX56抗体	
Alias:	61 kd nucleolar helicase; ATP-dependent 61 kDa nucleolar RNA helicase; DDX21; DDX26; Ddx56; DDX56_HUMAN; DEAD (Asp-Glu-Ala-Asp) box helicase 56; DEAD (Asp-Glu-Ala-Asp) box polypeptide 56; DEAD box protein 21; DEAD box protein 56; DEAD box RNA helicase; DEAD/H box 56; NOH61; Nucleolar helicase of 61 kDa; Probable ATP-dependent RNA helicase DDX56; Putative nucleolar RNA helicase; 8]	
Organism Species:	Rabbit	
Clonality:	Polyclonal	
React Species:	Human, Mouse, Rat, Cow, Horse, Sheep,	
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.	
Molecular weight:	62kDa	
Cellular localization:	The nucleus	
Form:	Lyophilized or Liquid	
Concentration:	1mg/ml	
immunogen:	KLH conjugated synthetic peptide derived from human DDX56:41-140/547	
Lsotype:	IgG	
Purification:	affinity purified by Protein A	
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.	
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.	
PubMed:	PubMed	
Product Detail:	This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA	

helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene shows ATPase activity in the presence of polynucleotides and associates with nucleoplasmic 65S preribosomal particles. This gene may be involved in ribosome synthesis, most likely during assembly of the large 60S ribosomal subunit. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012]

Function:

May play a role in later stages of the processing of the pre-ribosomal particles leading to mature 60S ribosomal subunits. Has intrinsic ATPase activity.

Subcellular Location:

Nucleus: nucleolus.

Tissue Specificity:

Detected in heart, brain, liver, pancreas, placenta and lung.

Post-translational modifications:

Phosphorylated upon DNA damage, probably by ATM or ATR.

Similarity:

Belongs to the DEAD box helicase family. DDX56/DBP9 subfamily.

Contains 1 helicase ATP-binding domain.

Contains 1 helicase C-terminal domain.

SWISS:

Q9NY93

Gene ID:

54606

Database links:

Entrez Gene: 54606 Human

Omim: 608023 Human

SwissProt: Q9NY93 Human

<u>Unigene: 654762</u> Human

Importa	nt Note:
---------	----------

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.suniondbiotech.com