



Rabbit Anti-EVI5L antibody

SL14650R

Product Name:	EVI5L
Chinese Name:	EVI5L蛋白抗体
Alias:	Ecotropic viral integration site 5 like protein; ecotropic viral integration site 5-like; EVI5L HUMAN; EVI5 like protein.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,Rabbit,Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	91kDa
Cellular localization:	The nucleocytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human EVI5L:701-794/794
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	EVI5L functions as a GTPase-activating protein (GAP) with a broad specificity. May interact with RAB10. Function: Functions as a GTPase-activating protein (GAP) with a broad specificity.

Subunit:

May interact with RAB10.

Subcellular Location:

Intracellular

Similarity:

Contains 1 Rab-GAP TBC domain.

SWISS:

Q96CN4

Gene ID:

115704

Database links:

[Entrez Gene: 115704](#) Human


[GenBank: NM_145245](#) Human

[SwissProt: Q96CN4](#) Human

[Unigene: 26870](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EVI5L) Polyclonal Antibody, Unconjugated (SL14650R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.