

Rabbit Anti-C6orf138 antibody

SL15220R

C6orf138 6号染色体开放阅读框138抗体 C6orf138; PTHD4_HUMAN; Patched domain-containing protein C6orf138. Rabbit Polyclonal Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit, WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-
C6orf138; PTHD4_HUMAN; Patched domain-containing protein C6orf138. Rabbit Polyclonal Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,
Rabbit Polyclonal Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,
Polyclonal Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit,
Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit,
WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-
500IF=1:100-500 (Paraffin sections need antigen repair)
not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.
96kDa
The cell membrane
Lyophilized or Liquid
1mg/ml
KLH conjugated synthetic peptide derived from human C6orf138:1-100/846
IgG
affinity purified by Protein A
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<u>PubMed</u>
Making up nearly 6% of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatibility complex proteins, which are key

molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf138 gene product has been provisionally designated C6orf138 pending further characterization.

Subcellular Location:

Membrane; Multi-pass membrane protein (Potential).

Similarity:

Belongs to the patched family.

Contains 1 SSD (sterol-sensing) domain.

SWISS:

Q6ZW05

Gene ID:

442213

Database links:

Entrez Gene: 442213Human

SwissProt: Q6ZW05Human

Unigene: 659409Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.