


## Rabbit Anti-SCOT2 antibody

## SL15589R

Product Name:	SCOT2
Chinese Name:	含氧酸辅酶A转移酶2A抗体
Alias:	SCOT2_HUMAN; Succinyl-CoA:3-ketoacid coenzyme A transferase 2, mitochondrial; 3-oxoacid CoA-transferase 2A; Testis-specific succinyl-CoA:3-oxoacid CoA-transferase; SCOT-t; OXCT2; FKSG25; SCOTT.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human, Mouse, Rat, Chicken, Horse, Rabbit, Sheep,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	52kDa
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human SCOT2:101-200/517
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	OXCT2 is a testis-specific succinyl-CoA:3-oxoacid CoA transferase (EC 2.8.3.5), which catalyzes the reversible transfer of CoA from succinyl-CoA to acetoacetate in the first step of ketone body utilization. See also OXCT1 (MIM 601424).[supplied by OMIM, Mar 2008].  Function:

Key enzyme for ketone body catabolism. Transfers the CoA moiety from succinate to acetoacetate. Formation of the enzyme-CoA intermediate proceeds via an unstable anhydride species formed between the carboxylate groups of the enzyme and substrate.
Subunit:
Homodimer.
Subcellular Location:
Mitochondrion.
Tissue Specificity:
Testis specific.
Similarity:
Belongs to the 3-oxoacid CoA-transferase family.
SWISS:
Q9BYC2
Gene ID:
64064
Gene ID: 64064 Database links:

Important Note:
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:


SY5Y(Human) Cell Lysate at 30 ug

Primary: Anti- SCOT2 (SL15589R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 52 kD

Observed band size: 52 kD



## Sample:

293T(Human) Cell Lysate at 30 ug

Primary: Anti- SCOT2 (SL15589R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 52 kD

Observed band size: 52 kD