



Rabbit Anti-VEGF-C antibody

SL1586R

Product Name:	VEGF-C
Chinese Name:	血管内皮生长因子C型抗体
Alias:	Vascuoar endothelial growth factor-C; AW228853; Flt4 ligand; Flt4-L; VEGF2; VEGFC; VRP; VEGFC_HUMAN.
文献引用 PubMed :	<p>Specific References(4) SL1586R has been referenced in 4 publications.</p> <p>[IF=7.84]Zhuo, Wei, et al. "The CXCL12?CCXCR4 Chemokine Pathway: A Novel Axis Regulates Lymphangiogenesis."Clinical Cancer Research 18.19 (2012): 5387-5398.Human, Mouse. PubMed:22932666</p> <p>[IF=5.62]He, Ting, et al. "Tumor cell-secreted angiogenin induces angiogenic activity of endothelial cells by suppressing miR-542-3p." Cancer Letters (2015).WB;Human. PubMed:26272182</p> <p>[IF=4.01]Alunno, Alessia, et al. "Mobilization of lymphatic endothelial precursor cells and lymphatic neovascularization in primary Sj?grens syndrome." Journal of Cellular and Molecular Medicine (2016).IHC-P;Human. PubMed:26828975</p> <p>[IF=0.72]Cialdai, Francesca, et al. "Modeled Microgravity Affects Fibroblast Functions Related to Wound Healing." Microgravity Science and Technology: 1-12.WB;Mouse. PubMed:0</p>
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections

	<p>need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.</p>
Molecular weight:	46kDa
Cellular localization:	Secretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human VEGF-C:321-415/415
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>Vascular endothelial growth factors (VEGFs), also known as vasculotropins, are a family of closely related growth factors having a conserved pattern of eight cysteine residues and sharing common VEGF receptors. VEGFs stimulate the proliferation of endothelial cells, induce angiogenesis, promote cell migration, increase vascular permeability, and inhibit apoptosis. The mitogenic activity of VEGFs appears to be mediated by specific VEGF receptors. The target cell specificity of VEGF is restricted to vascular endothelial cells. Vascular Endothelial Growth Factor C (VEGFC) is a member of the VEGF subfamily of PDGF-related growth factors. It is the ligand for Flt4 (VEGFR3) and KDR (VEGFR2). VEGFC binds Flt4 and induces tyrosine autophosphorylation of VEGFR3 and VEGFR2. VEGFC also stimulates the migration of bovine capillary endothelial cells in collagen gel. It is a specific growth factor for the lymphatic vascular system and mediates lymphangiogenesis. VEGFC is abundantly expressed in heart and skeletal muscle. Other tissues such as lung and kidney also express VEGFC.</p> <p>Subunit: Homodimer; non-covalent and antiparallel.</p> <p>Subcellular Location: Secreted.</p> <p>Tissue Specificity: Spleen, lymph node, thymus, appendix, bone marrow, heart, placenta, ovary, skeletal muscle, prostate, testis, colon and small intestine and fetal liver, lung and kidney, but not in peripheral blood lymphocyte.</p> <p>Similarity: Belongs to the PDGF/VEGF growth factor family.</p> <p>SWISS:</p>

P97953

Gene ID:
7424

Database links:

[Entrez Gene: 7424](#)Human

[Entrez Gene: 22341](#)Mouse

[Entrez Gene: 114111](#)Rat

[Omim: 601528](#)Human

[SwissProt: P49767](#)Human

[SwissProt: P97953](#)Mouse

[SwissProt: O35757](#)Rat

[Unigene: 435215](#)Human


[Unigene: 1402](#)Mouse

[Unigene: 6913](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Paraformaldehyde-fixed, paraffin embedded (Rat small intestine); Antigen retrieval by microwave in sodium citrate buffer (pH6.0) ; Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (VEGF-C) Polyclonal Antibody, Unconjugated (SL1586R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP)and DAB staining.

www.sunlongbiotech.com