



Rabbit Anti-phospho-ATP1A1 (Tyr260) antibody

SL15967R

Product Name:	phospho-ATP1A1 (Tyr260)
Chinese Name:	磷酸化钠钾ATP酶蛋白a1抗体
Alias:	alpha 1 Sodium Potassium ATPase (phospho Y260); alpha 1 Sodium Potassium ATPase; A1A1; AT1A1; AT1A1_HUMAN; Atpa-1; ATPase Na ⁺ /K ⁺ transporting alpha 1 polypeptide; ATPase Na ⁺ /K ⁺ transporting subunit alpha 1; BC010319; EC 3.6.3.9; MGC3285; MGC38419; MGC51750; Na K ATPase alpha A catalytic polypeptide; Na K ATPase catalytic subunit alpha A protein; Na ⁽⁺⁾ /K ⁽⁺⁾ ATPase 1; Na ⁽⁺⁾ /K ⁽⁺⁾ ATPase alpha-1 subunit; Na ⁺ , K ⁺ ATPase alpha subunit; Na ⁺ /K ⁺ ATPase alpha 1 subunit; Na ⁺ /K ⁺ ATPase 1; Na,K ATPase alpha 1 subunit; Nkaa1b; Sodium potassium ATPase alpha 1 polypeptide; Sodium pump 1; Sodium pump subunit alpha-1; sodium-potassium ATPase catalytic subunit alpha-1; Sodium/potassium-transporting ATPase subunit alpha-1.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,Horse,Sheep,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	113kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthesised phosphopeptide derived from human alpha 1 Sodium Potassium ATPase around the phosphorylation site of Tyr260:V(p-Y)TG
Isotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized

	antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>The protein encoded by this gene belongs to the family of P-type cation transport ATPases, and to the subfamily of Na⁺/K⁺-ATPases. Na⁺/K⁺ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The catalytic subunit of Na⁺/K⁺ -ATPase is encoded by multiple genes. This gene encodes an alpha 1 subunit. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May2009].</p> <p>Function: This is the catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. This action creates the electrochemical gradient of sodium and potassium ions, providing the energy for active transport of various nutrients.</p> <p>Subcellular Location: Cell membrane. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.</p> <p>Post-translational modifications: Phosphorylation on Tyr-10 modulates pumping activity.</p> <p>Similarity: Belongs to the cation transport ATPase (P-type) (TC 3.A.3) family. Type IIC subfamily.</p> <p>SWISS: P05023</p> <p>Gene ID: 476</p> <p>Database links:</p> <p>Entrez Gene: 476Human Entrez Gene: 11928Mouse Entrez Gene: 397481Pig Entrez Gene: 24211Rat Omim: 182310Human</p>

[SwissProt: P05023](#)Human

[SwissProt: Q8VDN2](#)Mouse

[SwissProt: P06685](#)Rat

[Unigene: 371889](#)Human

[Unigene: 280103](#)Mouse

[Unigene: 217534](#)Rat

[Unigene: 2992](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com