



Rabbit Anti-HOPX antibody

SL17356R

Product Name:	HOPX
Chinese Name:	绒毛膜癌不表达蛋白1抗体
Alias:	CAMEO; heart odd homeobox 1 protein; HOD; Homeodomain-only protein; HOP; HOP homeobox; HOP_HUMAN; HOPX; LAGY; Lung cancer-associated Y protein; NECC1; Not expressed in choriocarcinoma clone 1; Not expressed in choriocarcinoma protein 1; OB1; odd homeobox 1 protein; Odd homeobox protein 1; OTTHUMP00000158970; SMAP31; TOTO.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Pig,Cow,Horse,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	8kDa
Cellular localization:	The nucleus
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human HOPX:21-73/73
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	The protein encoded by this gene is a homeodomain protein that lacks certain conserved residues required for DNA binding. It was reported that choriocarcinoma cell lines and tissues failed to express this gene, which suggested the possible involvement of this

gene in malignant conversion of placental trophoblasts. Studies in mice suggest that this protein may interact with serum response factor (SRF) and modulate SRF-dependent cardiac-specific gene expression and cardiac development. Multiple alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, Feb 2009]

Function:

Atypical homeodomain protein which does not bind DNA and is required to modulate cardiac growth and development. Acts via its interaction with SRF, thereby modulating the expression of SRF-dependent cardiac-specific genes and cardiac development. Prevents SRF-dependent transcription either by inhibiting SRF binding to DNA or by recruiting histone deacetylase (HDAC) proteins that prevent transcription by SRF. Overexpression causes cardiac hypertrophy (By similarity). May act as a tumor suppressor.

Subcellular Location:

Nucleus.

Tissue Specificity:

Widely expressed. Expressed in the heart, brain, placenta, lung, skeletal and smooth muscles, uterus, urinary bladder, kidney and spleen. Down-regulated in some types of cancer such as lung cancer, choriocarcinoma, head and neck squamous cell carcinoma and oral squamous cell carcinoma.

Similarity:

Contains 1 homeobox DNA-binding domain.

SWISS:

Q9BPY8

Gene ID:

84525

Database links:

[Entrez Gene: 281270](#) Cow

[Entrez Gene: 84525](#) Human

[Entrez Gene: 74318](#) Mouse

[Entrez Gene: 396692](#) Pig

[Entrez Gene: 171160](#) Rat

[Omim: 607275](#) Human

[SwissProt: Q8MJD5](#) Cow

[SwissProt: Q9BPY8](#) Human

[SwissProt: Q8R1H0](#) Mouse

[SwissProt: Q8MJD6](#) Pig

[SwissProt: Q78ZR5](#) Rat

[Unigene: 619396](#) Human

[Unigene: 181852](#) Mouse

[Unigene: 478741](#) Mouse

[Unigene: 2989](#) Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com