

Rabbit Anti-MSL3L2 antibody

SL17859R

MSL3L2
MSL3L2蛋白抗体 A A A A A A A A A A A A A A A A A A A
Male specific lethal 3 homolog 2; Male specific lethal 3 like 2 (Drosophila); MSL3 like 2.
Rabbit
Polyclonal
Human,
WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100- 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
51kDa
The nucleus
Lyophilized or Liquid
1mg/ml
KLH conjugated synthetic peptide derived from human MSL3L2:1-100/447
IgG
affinity purified by Protein A
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed
MSL3L2 contains 1 chromo domain. It may be involved in chromatin remodeling and transcriptional regulation.Function: May be involved in chromatin remodeling and transcriptional regulation.

	Subcellular Location:
	Nuclear
	Similarity:
	Contains 1 chromo domain.
	Contains 1 MRG domain.
	SWISS:
	P0C860
	Gene ID:
	151507
	Database links:
	Database links: Entrez Gene: 151507 Human SwissProt: P0C860 Human
	SwissProt: P0C860 Human
	XO
	Important Note:
	This product as supplied is intended for research use only, not for use in human,
	therapeutic or diagnostic applications.
Picture:	×
	Sample:
	Hcclm3(Human) Cell Lysate at 30 ug
	Primary: Anti-MSL3L2 (SL17859R) at 1/300 dilution
	Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
	Predicted band size: 51 kD
	Observed band size: 51 kD