


Rabbit Anti-Melanoma Inhibitory Activity antibody

SL18783R

Melanoma Inhibitory Activity
黑色素瘤抑制活性蛋白抗体
Cartilage derived retinoic acid sensitive protein; CD RAP; CDRAP; Melanoma derived growth regulatory protein precursor; Melanoma inhibitory activity protein; Melanoma-derived growth regulatory protein; Mia; MIA_HUMAN.
Rabbit
Polyclonal
Human, Mouse, Rat, Pig, Cow, Horse, Rabbit, Sheep,
ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
12kDa
Secretory protein
Lyophilized or Liquid
1mg/ml
KLH conjugated synthetic peptide derived from human Melanoma Inhibitory Activity:21-100/131
IgG
affinity purified by Protein A
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<u>PubMed</u>



Picture:

Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MIA) Polyclonal Antibody, Unconjugated (SL18783R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.