



Rabbit Anti-ZFP62 antibody

SL19128R

Product Name:	ZFP62
Chinese Name:	Zinc finger protein62抗体
Alias:	DKFZp667F2013; FLJ11344; MGC176438; Zfp-62; Zfp62; ZFP62_HUMAN; Zinc finger protein 62 homolog; ZNF755.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Pig,Cow,Horse,Rabbit,Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	103kDa
Cellular localization:	The nucleus
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human ZFP62:431-530/900
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	Annotation of ZFP62 was placed on AC016572.6 on NCBI's annotation 36.3. This clone has been determined to be redundant, so the annotation was merged with the location on the assembled chromosome. [02 Mar 2009] Function: May play a role in differentiating skeletal muscle.

Subcellular Location:

Nucleus.

Similarity:

Belongs to the krueppel C2H2-type zinc-finger protein family.
Contains 23 C2H2-type zinc fingers.

SWISS:

Q8NB50

Gene ID:

643836

Database links:

[Entrez Gene: 643836](#) Human

[Entrez Gene: 22720](#) Mouse

[Oimim: 610281](#) Human


[SwissProt: Q8NB50](#) Human

[SwissProt: Q8C827](#) Mouse

[Unigene: 509227](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Paraformaldehyde-fixed, paraffin embedded (rat testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP62) Polyclonal Antibody, Unconjugated (SL19128R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.