


## Rabbit Anti-NME3 antibody

## SL19294R

Product Name:	NME3
Chinese Name:	二磷酸核苷激酶3抗体
Alias:	DR-nm23; Expressed in non metastatic cells 3; NDK 3; NDK3; NDK3_HUMAN; NDP kinase 3; NDP kinase C; NDPK C; NDPKC; NM23 H3; nm23-H3; NME 3; Nme3; Non metastatic cells 3 protein expressed in; Nucleoside diphosphate kinase 3; Nucleoside diphosphate kinase C.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human, Mouse, Rat, Pig, Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	19kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	lmg/ml
immunogen:	KLH conjugated synthetic peptide derived from human NME3:81-169/169
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	



## Picture:

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NME3) Polyclonal Antibody, Unconjugated (SL19294R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.