



Rabbit Anti-NUDT12 antibody

SL19509R

Product Name:	NUDT12
Chinese Name:	NUDT12蛋白抗体
Alias:	0610016O18Rik; DKFZP761I172; EC 3.6.1.22; Nucleoside diphosphate linked moiety X motif 12; Nucleoside diphosphate linked moiety X type motif 12; Nudix (nucleoside diphosphate linked moiety X) type motif 12; Nudix motif 12; Nudix type motif 12; NUDT 12; Peroxisomal NADH pyrophosphatase NUDT12.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,Rabbit,Sheep,
Applications:	WB=1:500-2000ELISA=1:500-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	52kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human NUDT12:201-300/462
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	Nucleotides are involved in numerous biochemical reactions and pathways within the cell as substrates, cofactors, and effectors. Nudix hydrolases, such as NUDT12, regulate the concentrations of individual nucleotides and of nucleotide ratios in response to changing circumstances (Abdelraheim et al., 2003 [PubMed 12790796]).[supplied by OMIM, Mar 2008]

Function:

Nucleotides are involved in numerous biochemical reactions and pathways within the cell as substrates, cofactors, and effectors. Nudix hydrolases, such as NUDT12 (Nudix (nucleoside diphosphate linked moiety X)-type motif 12), regulate the concentrations of individual nucleotides and of nucleotide ratios in response to changing circumstances. NUDT12 hydrolyzes NAD(P)H to NMNH and AMP (2',5'-ADP), and diadenosine diphosphate to AMP. It also has moderate activity towards NAD(P)(+), ADP-ribose and diadenosine triphosphate. It may regulate the concentration of peroxisomal nicotinamide nucleotide cofactors required for oxidative metabolism.

Subcellular Location:

Peroxisome

SWISS:

Q9BQG2

Gene ID:

82594

Database links:

[Entrez Gene: 83594](#) Human

[Omir: 609232](#) Human


[SwissProt: Q9BQG2](#) Human

[Unigene: 434289](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Sample:

Liver (Mouse) Lysate at 40 ug

Primary: Anti-NUDT12 (SL19509R) at 1/500 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 52 kD

Observed band size: 52 kD