

## Rabbit Anti-SLC16A10 antibody

## SL19797R

Chinese Name:溶质载体家族蛋白16成员A10抗体Alias:Aromatic amino acid transporter 1; MCT 10; MCT10; Monocarboxylate transpo MOT10_HUMAN; SLC16A10; Solute carrier family 16 member 10; T-type am acid transporter 1; TAT1.Organism Species:RabbitClonality:PolyclonalReact Species:Human,Mouse,Rat, WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.Molecular weight:55kDaCellular localization:The cell membraneForm:Lyophilized or Liquid	
Alias:  MOT10_HUMAN; SLC16A10; Solute carrier family 16 member 10; T-type am acid transporter 1; TAT1.  Organism Species:  Rabbit  Clonality:  Polyclonal  Human,Mouse,Rat,  WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight:  Cellular localization: The cell membrane  Form:  Lyophilized or Liquid	
Clonality:  React Species:  Human,Mouse,Rat,  WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight:  Cellular localization: The cell membrane Form: Lyophilized or Liquid	
React Species:  Human,Mouse,Rat,  WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 55kDa  Cellular localization: The cell membrane  Evorm: Lyophilized or Liquid	
Applications:  WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight:  Cellular localization: The cell membrane  Lyophilized or Liquid	
Applications:  500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 55kDa  Cellular localization: The cell membrane  Lyophilized or Liquid	
Cellular localization: The cell membrane  Form: Lyophilized or Liquid	:100-
Cellular localization: The cell membrane  Form: Lyophilized or Liquid	
Concentration: 1mg/ml	
immunogen: KLH conjugated synthetic peptide derived from human SLC16A10:421-515/515 <extracellular></extracellular>	
Lsotype: IgG	
Purification: affinity purified by Protein A	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.	
Storage:  Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent antibody the antibody is stable for at least two weeks at 2-4 °C.	n a year
PubMed: PubMed	
Product Detail:	