



Rabbit Anti-P2RX1 antibody

SL21358R

Product Name:	P2RX1
Chinese Name:	三磷酸腺苷受体受体P2X1抗体
Alias:	P2X1; ATP receptor; ATP receptor; H sapiens mRNA for ATP receptor; H.sapiens mRNA for ATP receptor; P2 RX1; P2RX1; P2RX1 protein; P2RX1_HUMAN; P2X purinoceptor 1; P2X receptor subunit 1; P2X1; P2X1 receptor; Purinergic receptor; Purinergic receptor P2X ligand gated ion channel 1; Purinergic receptor P2X ligand-gated ion channel 1; Purinergic receptor P2X1.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Cow,Rabbit,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	45kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human P2RX1:
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	The P2X receptor family is comprised of ligand-gated ion channels that allow for the increased permeability of calcium into the cell in response to extracellular ATP. The seven P2X receptors, P2X1-P2X7, form either homomeric or heteromeric channels or

both. They are characterized by intracellular amino- and carboxy-termini. P2X receptors are expressed in a wide variety of tissues, including neurons, prostate, bladder, pancreas, colon, testis and ovary. The major function of the P2X receptors is to mediate synaptic transmissions between neurons and to other tissues via the binding of extracellular ATP, which acts as a neurotransmitter. The P2X receptors may be involved in the onset of necrosis or apoptosis after prolonged exposure to high concentrations of extracellular ATP.

Function:

Ligand-gated ion channel with relatively high calcium permeability. Binding to ATP mediates synaptic transmission between neurons and from neurons to smooth muscle. Seems to be linked to apoptosis, by increasing the intracellular concentration of calcium in the presence of ATP, leading to programmed cell death.

Subunit:

Homo- or heteropolymers (By similarity).

Subcellular Location:

Membrane; Multi-pass membrane protein.

Similarity:

Belongs to the P2X receptor family.

SWISS:

P51575

Gene ID:

5023

Database links:

[Entrez Gene: 5023](#)Human

[Entrez Gene: 18436](#)Mouse

[Entrez Gene: 25505](#)Rat

[Omim: 600845](#)Human

[SwissProt: P51575](#)Human

[SwissProt: P51576](#)Mouse

[SwissProt: P47824](#)Rat

[Unigene: 41735](#)Human


[Unigene: 25722](#)Mouse

[Unigene: 91176](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Sample:

Urinary bladder (Mouse) Lysate at 40 ug

Penis (Mouse) Lysate at 40 ug

Primary: Anti- P2RX1 (SL21358R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 45 kD

Observed band size: 65 kD

www.sunlongbiotech.com