



## Rabbit Anti-Phospho-MKP1 (Ser296) antibody

SL23272R

<b>Product Name:</b>	Phospho-MKP1 (Ser296)
<b>Chinese Name:</b>	磷酸化丝裂原活化蛋白激酶磷酸酶-1抗体
<b>Alias:</b>	DUSP1 (phospho S296); MKP1 (Phospho Ser296); MKP1 (Phospho S296); Mitogen activated protein kinase phosphatase 1; CL 100; CL100; Dual Specificity Phosphatase 1; Dual specificity protein phosphatase 1; Dual specificity protein phosphatase hVH1; DUSP 1; EC 3.1.3.16; EC 3.1.3.48; HVH 1; HVH1; MAP kinase phosphatase 1; MKP 1; MKP1; MKP-1; Protein tyrosine phosphatase CL100; PTPN 10; PTPN10; Serine/threonine specific protein phosphatase; VH 1; VH1; 3ch134; Mkp1; Ptpn16.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Pig,Cow,Horse,Rabbit,
<b>Applications:</b>	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	39kDa
<b>Cellular localization:</b>	cytoplasmic
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated Synthesised monomethylpeptide derived from human MKP1 around the phosphorylation site of Ser296:II(p-S)PN
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>

The expression of DUSP1 gene is induced in human skin fibroblasts by oxidative/heat stress and growth factors. It specifies a protein with structural features similar to members of the non-receptor-type protein-tyrosine phosphatase family, and which has significant amino-acid sequence similarity to a Tyr/Ser-protein phosphatase encoded by the late gene H1 of vaccinia virus. The bacterially expressed and purified DUSP1 protein has intrinsic phosphatase activity, and specifically inactivates mitogen-activated protein (MAP) kinase in vitro by the concomitant dephosphorylation of both its phosphothreonine and phosphotyrosine residues.

**Function:**

Dual specificity phosphatase that dephosphorylates MAP kinase MAPK1/ERK2 on both 'Thr-183' and 'Tyr-185'.

**Tissue Specificity:**

Expressed at high levels in the lung, liver placenta and pancreas. Moderate levels seen in the heart and skeletal muscle. Lower levels found in the brain and kidney.

**Similarity:**

Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.

Contains 1 rhodanese domain.

Contains 1 tyrosine-protein phosphatase domain.

**Product Detail:**

**SWISS:**

P28562

**Gene ID:**

1843

**Database links:**

[Entrez Gene: 1843](#)Human

[Entrez Gene: 19252](#)Mouse

[Entrez Gene: 114856](#)Rat

[Omim: 600714](#)Human

[SwissProt: P28562](#)Human

[SwissProt: P28563](#)Mouse

[SwissProt: Q64623](#)Rat

[Unigene: 171695](#)Human

[Unigene: 239041](#)Mouse

[Unigene: 98260](#)Rat

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

MKPs是一类丝氨酸/苏氨酸和酪氨酸双重底物特异性的磷酸酶,对于丝裂素活化蛋白激酶活性的调节起着十分重要的作用,可使丝裂素活化蛋白激酶上的苏氨酸/酪氨酸去磷酸化失活。目前研究发现MKPs分别有MKP-1、MKP-2、MKP-3及MKP4-6。

MKPs受MAPK信号通路中多种成分的诱导,决定了它与MAPK之间作用的特异性。

通过去磷酸化作用调节MAPK信号途径的活性,确保了细胞内信号的精确传递,参与了多种主要的细胞功能的调节。

www.sunlongbiotech.com