



Rabbit Anti-CCL1 precursor antibody

SL2450R

Product Name:	CCL1 precursor
Chinese Name:	嗜酸粒细胞趋化蛋白CCL1前体蛋白抗体
Alias:	I-309; TCA3; C-C motif chemokine 1 precursor; chemokine (C-C motif) ligand 1; BF534335; Scya1; Chemokine CC Motif Ligand 1; inflammatory cytokine I-309; P500; SCYA1; SISE; small inducible cytokine A1; Ccl1; Small-inducible cytokine A1; T lymphocyte secreted protein I-309; CCL1_HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Pig,Rabbit,
Applications:	ELISA=1:500-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	10kDa
Cellular localization:	Secretory protein
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human C-C motif chemokine 1 precursor:21-30/96
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two

cysteines separated by a single amino acid. This cytokine is secreted by activated T cells and displays chemotactic activity for monocytes but not for neutrophils. It binds to the chemokine receptor CCR8. [provided by RefSeq, Jul 2008].

Function:

Cytokine that is chemotactic for monocytes but not for neutrophils. Binds to CCR8.

Subunit:

Monomer.

Subcellular Location:

Secreted.

Similarity:

Belongs to the intercrine beta (chemokine CC) family.

SWISS:

P22362

Gene ID:

6346

Database links:

[Entrez Gene: 6346](#)Human

[Omim: 182281](#)Human

[SwissProt: P22362](#)Human

[Unigene: 72918](#)Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

CCL1属于CCChemokine家族, 是由活化的T细胞分泌的glycoprotein。CCL1是一小分子的cell factorC与细胞表面的Chemokine受体CCR8结合,对单核细胞, Natural killer cells, 未成熟的B细胞和树突状细胞有细胞趋化作用。