



## Rabbit Anti-GRO gamma antibody

SL2547R

<b>Product Name:</b>	GRO gamma
<b>Chinese Name:</b>	生长调节致癌基因gamma/GRO $\gamma$ 抗体
<b>Alias:</b>	growth $\square$ regulated oncogene gamma; Chemokine (CXC motif) ligand 3; Cinc 2; Cinc2; CINC2b; CXCL 3; CXCL3; Cytokine induced neutrophil chemoattractant 2; Dcip1; Dendritic cell inflammatory protein 1; Gm1960; GRO3; GRO3 oncogene; GROG; Growth regulated protein gamma; Macrophage inflammatory protein 2 beta precursor; Melanoma growth stimulatory activity gamma; MGSA gamma; MIP 2b; MIP2B; SCYB3; CXCL3 HUMAN.
<b>文献引用</b> <b>PubMed</b> :	<p><b>Specific References(1)</b> SL2547R has been referenced in 1 publications.</p> <p><b>[IF=3.53]</b>Gui S, Ni S, Jia J, Gong Y, Gao L, et al. (2014) Inconformity of CXCL3 Plasma Level and Placenta Expression in Preeclampsia and Its Effect on Trophoblast Viability and Invasion. PLoS ONE 9(12): e114408. <b>IHC-P;Human.</b></p> <p style="text-align: right;"><a href="#">PubMed:25485631</a></p>
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Pig,Cow,Horse,Rabbit,Sheep,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	12kDa
<b>Cellular localization:</b>	Secretory protein
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human GRO gamma:35-107/107
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A

<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	<p>Growth Regulated Protein gamma (GRO gamma) is a cytokine that may play a role in inflammation and affects endothelial cells in an autocrine fashion. It belongs to the intercrine alpha family (small cytokine c-x-c) (chemokine cxc).</p> <p><b>Function:</b> Ligand for CXCR2. Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.</p> <p><b>Subcellular Location:</b> Secreted.</p> <p><b>Post-translational modifications:</b> N-terminal processed form GRO-gamma(5-73) is produced by proteolytic cleavage after secretion from peripheral blood monocytes.</p> <p><b>Similarity:</b> Belongs to the intercrine alpha (chemokine CxC) family.</p> <p><b>SWISS:</b> P19876</p> <p><b>Gene ID:</b> 2921</p> <p><b>Database links:</b>  <a href="#">Entrez Gene: 2921</a>Human  <a href="#">Omim: 139111</a>Human  <a href="#">SwissProt: P19876</a>Human  <a href="#">Unigene: 89690</a>Human</p> <p><b>Important Note:</b> This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p>

黑素瘤生长刺激因子3.

[www.sunlongbiotech.com](http://www.sunlongbiotech.com)